



An update on biological advancement of *Jatropha curcas* L.: New insight and challenges

Purabi Mazumdar^{a,*}, Pooja Singh^a, Subramanian Babu^b, Ramamoorthy Siva^b, Jennifer Ann Harikrishna^{a,c}

^a Centre for Research in Biotechnology for Agriculture, University of Malaya, 50603 Kuala Lumpur, Malaysia

^b School of Bio Sciences and Technology, VIT University, 632014 Vellore, Tamil Nadu, India

^c Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

ARTICLE INFO

Keywords:

Jatropha

Biodiesel

Flower ratio

Abiotic and biotic stress

TAG biosynthesis

Transcription factors

ABSTRACT

Sustainable production of non-food oleaginous crops is highly desirable towards meeting increasing global demands for energy by supplementing the diminishing supply of fossil fuels without affecting food security. *Jatropha curcas* L., a high yielding oleaginous seed crop (30–48% oil), with a simple propagation system, short gestation period and high adaptability to a wide range of climatic and edaphic conditions, qualifies to be a potential candidate for sustainable biofuel production. The crop has been proposed as part of the solution to the challenges of climate change, energy scarcity and provision of rural income, as it can grow in marginal soils with minimal nutrient content. However, in 2010, this view was challenged due to unpredictable yield patterns of the crop, recorded for plantation in marginal and low nutrient environments. The main reason behind the failure to meet its promise, was the paucity of information of the optimal agricultural practices and nutrient requirements required for large scale cultivation of *J. curcas*. The low productivity of seed in *J. curcas* was identified to result from a combination of variation in seed quality and quantity, high male to female flower ratio, asynchronous flowering, seed toxicity and susceptibility to various biotic and abiotic stresses. With the advancement of new technology and the availability of a whole genome sequence for *J. curcas*, several studies addressed at overcoming the shortcomings of the plant via biological approaches have been reported, indicating a renewed promise for this energy crop. This review addresses recent developments in improvement of *J. curcas* from the biological perspective, identifies some remaining gaps in knowledge and provides recommendations for future studies towards economically viable *J. curcas* cultivation.

1. Introduction

Jatropha curcas L., belongs to the Euphorbiaceae family and is known for its multiple beneficial applications from biodiesel production [1] to the herbal medicine [reviewed in 2] and cosmetic industries [reviewed in 3]. *J. curcas* is indigenous to Mexico and Central America and has spread widely throughout the tropics [4]. The plant garnered worldwide attention as an alternative source of sustainable energy mainly because of its non-edible high seed oil content. Each mature plant of *J. curcas* produces an average of 4–5 kg of seeds per year when cultivated under suitable conditions with an extensive productive period of around 30–50 years [5]. In addition to its easy propagation by grafting or seed [6], short gestation period, high adaptability to a wide range of environments and soils [reviewed in 7], ability to sequester carbon in degraded soils [8] and capacity for the bioremediation of heavy metal-contaminated soils [9] makes *J. curcas* a crop of choice for

biodiesel production [10] in developing countries. As a non-food crop, capable of growth on land unsuitable for food production, and with good potential to address issues related to climate change, growing energy demand and the need to improve rural income, *J. curcas* was seen as an ideal energy crop. Thus in 2009, initiatives were taken by the governments of India, China and Brazil along with several biofuel companies and NGOs for large scale cultivation of *J. curcas* [11]. Other countries such as Myanmar, Malaysia and Malawi also showed their interest to grow this plant. However, in 2010, questions on the sustainability of Jatropha-based biofuel programs were raised, because of its underperformance in large scale cultivation [12–14]. The main reasons behind the failure of *J. curcas* to meet its promises were inadequacy in field validation and agricultural practices, lack of knowledge on nutritional requirements of the crop [15], high variation in seed quantity and quality, high male to female flower ratio [16], seed toxicity [17] and susceptibility to biotic [18] and abiotic stresses [19].

* Corresponding author.

E-mail address: purabi@um.edu.my (P. Mazumdar).

<https://doi.org/10.1016/j.rser.2018.04.082>

Received 21 February 2017; Received in revised form 22 June 2017; Accepted 14 April 2018
1364-0321/ © 2018 Elsevier Ltd. All rights reserved.

Subsequently, research on the biological improvement of *J. curcas* has been reported to address many of the apparent shortcomings of this crop.

Previous reviews have highlighted the multiple applications of *J. curcas* [20–26] such as use for biodiesel production [27–33], the status of *J. curcas* cultivation [34,35], socio-economic and environmental aspects [36–38]; sustainability issues [39], breeding aspects [40–42], tissue culture and transformation [7,41–44] biochemical studies [44], seed toxicity [21,26,45,46], usage of latest technology to analyse metabolic pathways [47] and various limitations of *J. curcas* in field conditions, from which suitable breeding strategies have been proposed for the future utilization of this plant [42,48,49]. In the present review, we examine recent studies conducted to address the limitations of *J. curcas* from the biological aspects of harnessing the knowledge on plant-microbe interactions, gene resources and biotechnological tools. We identify the gaps in biological knowledge and applications that remain for this crop and provide recommendations for future biological research for *J. curcas* towards economic feasibility for bioenergy production.

2. Review of studies using biological approaches to address the limitations of *J. curcas* as a biofuel plant

Some of the major challenges to adopt *J. curcas* as potential feedstock of future sustainable fuel have been summarised in Fig. 1. There are several biological approaches that can be used to address these challenges, as discussed in the following sections:

2.1. Optimising *J. curcas* productivity in Low nutrient soil

Poor productivity in low nutrient soil was reported as a major limitation to *J. curcas*'s success as a biofuel crop [4]. To avoid the competition for land use for food crops, the ideal for biodiesel crops is to be grown in low nutrient or barren soils. Due to high phenotypic plasticity, *J. curcas* can easily grow in such soil, but low nutrient soil reduces seed yield [15]. To improve the seed yield, nitrogen fertilizers were initially tested, which enhanced *J. curcas* growth and productivity [50]. However, emission of reactive N_2O from nitrogen fertilizers traps heat in the atmosphere 310 times more effectively than CO_2 [51] and hence is considered as one of the important contributors to global warming. To enrich the fertility level in barren and degraded land, organic amendments (such as animal manure) were used to evaluate germination and growth patterns of *J. curcas* in a field trial [15]. The field trial showed significant enhancement of *J. curcas* growth and biomass compared to the control plants. A disadvantage of the

application of organic amendments was that this attracted humivorous termites, which lead to severe damage of the seedlings [15]. To overcome the termite manifestation, use of pesticides has been recommended [15,52]. However, pesticide use risks encouraging the emergence of resistant pest in addition to environmental toxicity [53]. As an alternative, several groups of researchers investigated the effects of beneficial plant-microbe interactions mediated by arbuscular mycorrhizal fungi (AMF), endophytes and plant growth promoting rhizobacteria (PGPR) in *J. curcas*.

AMF transfer inorganic soil nutrients P, Zn and N to the host plant in exchange for organic carbon [54]. Such symbiotic associations of AMF with crop plants are widely known [reviewed in 55]. AMF diversity in the rhizosphere of *J. curcas* was explored in plantation areas in Thailand [56] and India [57,58] from which AMF were seen to promote *J. curcas* growth under a wide range of field conditions, from low to high nutrient soils including acidic, saline and alkaline conditions. Most common fungi species recorded in such symbiosis belong to the genera *Glomus*, *Acaulospora*, *Gigaspora* and *Scutellospora* [59,60]. Low nutrient soil (low P availability) supplemented with AMF spores (*Glomus aggregatum*, *G. fasciculatum*, *G. intraradices*, and *G. mosseae*) showed growth enhancement of *J. curcas* seedlings compared to non-supplemented soil [61]. Similarly, in a field trial, *J. curcas* inoculated with AMF spores (*Glomus fasciculatum* and *Scutellospora calospora*), showed enhanced plant height, higher branch number and larger canopy area compared to the plants supplied with NPK fertilizer [59]. *J. curcas* inoculated with AMF (*Glomus mosseae*, *G. microcarpum*, *G. fasciculatum*, *G. intraradices*, *Gigaspora margarita*, and *G. heterogama*) isolated from the rhizosphere of castor (another oilseed crop of the Euphorbiaceae family) showed improved plant vigor under salt stress conditions, suggesting a vital role of AMF in plant salinity-tolerance [57]. AMF (*Glomus*, *Scutellospora*, *Entrophosphora* and *Gigaspora*) inoculation also increased the survival rate of *J. curcas* grown in highly acidic, low nutrient soil [62]. *J. curcas* inoculated with AMF grown in heavy metal contaminated soil (Cu and Pb) was also reported to show reduced translocation of heavy metals from the roots to the aerial plant parts [63]. Hence, AMF can be successfully applied as a component of bio-fertilizers to counteract problems of nutrient deficiency, salinity and metal toxicity in soil. However, since mycorrhizal species often have specific ecological and edaphic requirements for efficient root colonization and plant growth promoting activity, such as soil pH, temperature and organic matter [reviewed in 64], the best performance of AMF in large-scale plantation, may be obtained by use of species indigenous to the particular edaphic condition.

Endophytes are mostly fungi and bacteria that live within living plant tissue. Several novel and beneficial endophytes identified from *J.*




Fig. 1. Limitation of *Jatropha curcas* as biodiesel plant.

Table 1Plant growth promoting endophytes reported in *J. curcas*.

Endophytes	Plant part	Beneficial features of Endophytes	References
Rhizobacteria	Roots	Promoted root and shoot growth	[65]
Enterobacter sp.	Roots	Stimulated plant growth and increased average seed number and seed weight in <i>J. curcas</i>	[66]
Actinobacteria (Streptomyces, Nonomuraea, Micrococcus and Kibdelosporangium sp.)	Roots, leaves, seeds and stems	Increased fresh weight, shoot and root length and leaf area of <i>J. curcas</i> seedlings	[67]
Methylobacterium sp.	Leaves	Promoted biomass and seed production in <i>J. curcas</i>	[68]
Firmicutes and Alphaproteobacteria (Paenibacillus, Brevibacillus, Sphingomonas and Rhizobium sp.)	Roots	Promoted growth and development of <i>Zea mays</i>	[69]

curcas, showed successful colonization in roots and enhanced plant growth and seed yield, even when grown in low nutrient soil [65–69] (see Table 1). Madhaiyan et al. [65] explored endophytic diversities in leaf, stem and root of *J. curcas*. Molecular and functional characterization of endophytes showed them to be species mainly belonging to α , β , γ -Proteobacteria, Actinobacteria, Firmicutes and Methylobacterium [68]. Among the Methylobacteria isolates, Methylobacterium L2–4 strain was characterized in detail and showed high nitrogen-fixing efficiency, and promoted biomass and seed production in *J. curcas*. Mohanty et al. [69] showed that endophytes isolated from *J. curcas* were also able to improve growth performance of other agricultural crops like maize.

In addition to endophytic bacteria, several PGPR have been identified with potential to enhance the growth and productivity of *J. curcas* on nutrient depleted or contaminated soils. Enterobacter sp. isolated from the rhizosphere of *J. curcas* grown in alkaline, wasteland, sodic and saline soils were shown to be efficient phosphate solubilizers and to have auxin production capacity during screening [70,71], however, their effectiveness in field conditions has not yet been demonstrated. A Chromobacterium sp. isolated as a PGRP from the rhizosphere of *J. curcas*, displayed an ability to degrade dichlorobenzene, a ground water pollutant [72]. Some PGPR sp. that have been commonly used for other crops, such as *Enterobacter cloacae*, *Pseudomonas pseudoalcaligenes* [73], *Pseudomonas fluorescens*, *Pseudomonas putida* [74] and *Bacillus* sp. [73,75] are also able to colonize the rhizosphere of *J. curcas* and promote fast seed germination and plant growth. While there has been advancement in knowledge of beneficial microbes for yield and growth improvement of *J. curcas* and that will potentially address issues of low productivity in low nutrient and marginal soils, further studies are still needed both for field scale evaluation and towards the low-cost mass production and high storage viability of microbial inoculants for sustainable cultivation.

In addition to soil enrichment, use of cultivars with improved seed yield and stress tolerance capacity for large scale plantation can enhance the productivity of *J. curcas* without effecting agricultural land use. Reviews have summarised promising research outcomes from morphological and molecular screening of elite accessions of *J. curcas* [reviewed in 7, 41]. A requisite for genetic improvement through breeding is an availability of sufficient genetic variability. Studies on morphometric traits such as plant growth, architecture, seed yield, seed weight and oil content among *J. curcas* accessions have shown the existence of considerable variability. However, diversity assessment based solely on morphometric traits are not always reliable, as *J. curcas* shows high plasticity. Additional information from studies with molecular markers including random amplified polymorphic DNA (RAPD), inter simple sequence repeats (SSR), amplified fragment length polymorphism (AFLP) and sequence-tagged sites (STS) suggest low genetic diversity except among accessions collected from central America and Mexico [76]. Thus, accessions identified from central America and Mexico show the best potential for use in breeding programs [77]. Several research groups worldwide viz, Wageningen University and Research Centre, Plant Research International (Netherlands) and Lao Institute of Renewable Energy (Laos), Surfactant and Bio-energy

Resource Centre (Indonesia), Agricultural Research Trust (Zimbabwe), National Oilseeds and Vegetable Oils Development Board of India and Indian Institute of Oilseeds Research (India) have collected and maintain *J. curcas* germplasm repository, housing several thousand accessions of *J. curcas* from various locations around the world. Many of the accessions have been screened for seed and oil traits towards support for breeding and development of stable elite varieties [reviewed in 7]. The details of selection strategies and breeding status of *J. curcas* in Brazil were discussed in de Azevedo et al. [78]. However, release of commercial elite cultivars via breeding is still in its infancy. As *J. curcas* is a perennial species with several traits of interest (as described in Fig. 1), the procedure of selection via conventional breeding is expected to be very slow (approximately 10–14 years) [78]. Several efficient transformation systems [79,80] and genetically modified *J. curcas* with improved oil trait [81], stress tolerance [82,83] have been reported and can be used in crop improvement. As for any crop, application of a combinatorial approach encompassing biotechnological tools and conventional breeding, are pivotal to expedite the development and release of elite cultivars.

2.2. Enhancing seed yield by increase in female flower ratio

One of the major shortcomings of *J. curcas* for biofuel production, is the unpredictable seed yield, with low seed production being a barrier to commercial success. *J. curcas* plants are mainly monoecious with flowers arranged as an axillary panicle polychasial cymes inflorescence (Fig. 2a), where the female flower is present at the top surrounded by sub branches which mainly produce male flowers (Fig. 2b). Occasionally hermaphrodite flowers are also observed in *J. curcas* [84,85]. *J. curcas* has a complex floral biology, with floral development divided into 12 phases [86]. The female flower develops as a hermaphrodite up to the sixth phase of floral development and thereafter sexual differentiation begins when stamens are aborted and the pistil develops [86]. In each inflorescence, the male to female ratio has been variously reported as 13.4:1 [87], 27: 1 [88] and 30:1 [16]. The variability in male and female flower ratio and relatively low number of female flowers is one of the reasons for variation of seed yield in *J. curcas* [89].

In recent years, several approaches have been explored towards increasing the female flower proportion in *J. curcas* inflorescences. Exogenous application of plant growth regulators (PGRs) such as gibberellin [90], cytokinin [91], paclobutrazol [92–94], auxin [95] and thidiazuron (TDZ) [96] to the inflorescence showed induction of larger numbers of bisexual flowers, increased female flower number and a 5–11-fold increase in *J. curcas* seed yield. These studies indicated great potential for improvement in *J. curcas* seed yield by the application of PGRs via significant increase in female flower ratio to increase fruit production and hence seed yield. Nevertheless, the application of PGR directly to flowers is labour intensive and not an economical and practical solution for a large-scale commercial plantation. Manipulation of genetic control of flowering may provide a better and long-term solution. Several key floral regulatory network genes have been identified in *J. curcas*: Li et al. [97] isolated *J. curcas* flowering locus T




Fig. 2. *Jatropha curcas* inflorescence (a) *J. curcas* plant with flowers arranged in axillary panicle polychasial cyme inflorescences. (b) Male flowers (M) and female flowers (F).

homolog (*JcFT*), which was observed to be expressed mainly in the female flowers, fruits, and seeds. Over-expression of *JcFT* in *J. curcas* resulted in early flowering transgenic lines supporting its role in inducing flower development [97]. Another flower development associated gene *LEAFY* (*JcLFY*) isolated from *J. curcas*, was shown to have a critical role in the early stage of flower bud formation, as over-expression induced early flowering, solitary flowers, and terminal flowers in *J. curcas* [98]. From a study comparing relative gene expression in tissues associated with flower development in a superior genotype of *Jatropha* (with high female flower number) and a genotype with low relative numbers of female flowers, abundant transcripts of *SUP* (Superman), *TAA1* (L-tryptophan-pyruvate aminotransferase 1), *CRY2* (Cryptochrome-2), and *CKX1* (Cytokinin dehydrogenase 1) were associated with increased numbers of female flowers [16]. This was explained based on the role of these genes in arresting the development of stamens [16]. Another study which profiled gene expression of *J. curcas* flower buds at six different developmental stages starting from prior to sex differentiation stages and up to mature female flower formation, provided intensive information on genes involved in carpel and stamen development and maturation of male and female flowers [99]. Among 15 useful sex-related candidate genes, gibberellin-regulated protein 4-like protein, *CLAVATA1* and adenosine monophosphate-activated protein kinase were associated with stamen differentiation; auxin response factors 6-like protein, *AGAMOUS*-like 20, *RING-H2* finger protein *ATL3J*, auxin-induced protein 22D, *CLAVATA1*, and *R2R3-MYB* transcription factor were linked to embryo sac development, while cytokinin oxidase, auxin repressed-like protein1-like protein, Gibberellin receptor protein 1 and auxin-induced protein X10A were present both during stamen differentiation and embryo sac development [99]. While further detailed functional validation is still needed, the identification of candidate genes from these types of study provides a good foundation for the future manipulation of floral ratio in *J. curcas*.

In addition to the importance of the female flower ratio in inflorescences, efficient pollination, aided by the establishment of pollination services in large scale plantations, aids to maximize fruit yield in *J. curcas* [reviewed in 100]. The studies on floral structure and pollination ecology of *J. curcas* accessions in Thailand [101], India [87,102–104], Brazil [105], China [106,107], Indonesia [108] and Mexico [109] showed that its flower exhibits protandry (in which male reproductive organs mature before the female ones) which favours xenogamy (cross pollination) over geitonogamy (self-pollination). Fruit production was recorded to be highest in open-pollination and in xenogamy. The pollinators of *J. curcas* mainly belong to the orders Hymenoptera, Lepidoptera and Diptera [108]. Studies conducted in different geographical locations showed honey bees (*Apis florea*, *A. dorsata*, *A. mellifera*, *Scaptotrigona mexicana* and *Trigona angustula*) [102,103,108–110] to be the most efficient pollinators compared to other insect visitors. This information can be strategically used to increase seed yield by establishment of honey bee hubs along with native pollinators in large scale plantation.

2.3. Detoxification of *J. curcas* seed cake using microbes

J. curcas seed cake is a by-product of oil extraction with high nutritional content, but also containing several non-desired compounds. The seed cakes have higher nutritional quality than soybean meal [111] with high levels of protein (45.3–58.6%), dietary fibre (5–5.5%) and with a gross energy value of 19–48% [112]. Essential amino acid profiling of *J. curcas* seed cake showed higher levels of valine, isoleucine, leucine, histidine and threonine than the recommended dosage for children by FAO/WHO [112]. The rich nutrient content would be ideal for human and animal nutrition, as fertilizer and as a substrate for production of industrial enzymes such as protease and lipase, as it supports good bacterial growth [113–115]. However, the presence of toxins such as phorbol esters, curcins and antinutritional compounds such as phytic acid, tannins and saponin make the seed cake not suitable as food or animal feed currently [112,116]. The use of *J. curcas* seed cake as a biofertilizer is also limited due to the phytotoxic properties of curcins [117–119]. Phorbol esters are highly toxic and carcinogenic, inducing leakage of plasma, tumors and tissue damage in humans and animals [120]. Curcins, has also been reported as stimuli for mucosal irritation and gastrointestinal hemagglutination in monogastric animals and humans [121]. The seed cake component phytic acid hinders the absorption of minerals in the intestines [122] while tannins reduce protein digestibility and stimulate liver damage [123]. Thus, the use of seed cake as an animal feed or fertilizer will only become feasible if the cake can be detoxified, which will not only aid commercial value to this by-product but would also diminish the environmental hazards caused by its inappropriate disposal.

In earlier studies, physical methods including heat-treatment, were able to detoxify curcins and tannins to a great extent, but not phorbol ester [124]. Among the toxins, phorbol esters have been recorded as heat resistant and the key toxic agents in *J. curcas* seed cake [116], where concentrations are reported to range from 0.21 to 0.47 mg/g [125]. To detoxify phorbol ester, chemical methods such as solvent extraction [17,21,111,126–131] and alkali treatments [132] have been developed and have shown up to 98% toxin reduction efficiency. However, a drawback of using chemical methods for detoxification are that the chemicals introduced in the process may result in other chemical residues. The progress on physical, chemical and biological detoxification processes for *J. curcas* seed cake was reviewed in Makkar et al. [133]. Efficiency and environmental safety are major advantages of biological detoxification over physical and chemical methods and recent research in detoxification has focused on eco-friendly biological detoxification using microorganism-mediated fermentation, and on toxin-free seed production using biotechnological tools. Biological detoxification is carried out using various strains of fungi and bacteria, several of which are efficient in toxin-reduction, as summarised in Table 2. The efficiency of Basidiomycete (Polyporales, Agaricales) and ascomycete (Saccharomycetales, Mucorales, Eurotiales and Hypocreales) fungi in detoxification of seed cake has been demonstrated in

Table 2
J. curcas seed cake detoxification experiments conducted by using biological agents.

Organism used	Experimental conditions for seed cake incubation with organisms	Targeted Toxin	Reduction (%)	Seed cake tested in animal feeding trial	Results of animal feeding trial	References
<i>Bjerkandera adusta</i>	Incubation at 28 °C for 30 days	Phorbol ester	91%	-	-	[198]
<i>Phelbia rufa</i>			97%	-	-	
<i>Ganoderma resinacum</i>			No significant reduction	-	-	
<i>Aspergillus niger</i>	Incubation for 7 days followed by growth termination by oven drying at 70 °C	-	-	African dwarf goats were fed for 112 days	Seed cakes promoted tissue development in goats	[199]
<i>Penicillium chrysogenum</i>		-	-			
<i>Trichoderma longibrachiatum</i>		-	-			
<i>Pseudomonas aeruginosa</i> PseA	Incubation at 30 °C, pH 7.0 and relative humidity 65% for 9 days	Phorbol esters	100%	-	-	[135]
<i>Lactobacillus plantarum</i>	Ensiling (60 days) by adding soluble carbohydrates in the cake and inoculant	Phorbol esters	47.40%	-	-	[136]
<i>Propionibacterium</i>						
<i>Pleurotus ostreatus</i>	Incubated at 28 °C for 30 days	Phorbol ester	99%	Female alpine goats were fed for 60 days	No toxic effect on animals	[134]
		Tannin	46%			
		Phytic acid	95%			
<i>Bacillus licheniformis</i>	Solid-state fermentation at 30 °C (for <i>B. subtilis</i>) and 37 °C (for <i>B. licheniformis</i>) for 7 days	Phorbol ester	Phorbol esters (32.8%), Phytic acid (57%), Trypsin inhibitor (1.2%)	-	-	[137]
<i>Bacillus subtilis</i>		Phytic acid Trypsin inhibitor				
<i>Aspergillus oryzae</i>	Solid-state fermentation for 12 days	Phorbol ester Phytic acid	Phorbol ester (57.8%) Phytic acid (92.9%)	-	-	[200]
<i>Aspergillus terreus</i>			Phorbol ester (57.8%) Phytic acid (92.1%)	-	-	
<i>Aspergillus versicolor</i>			Phorbol ester (68.6%) Phytic acid (64.6%)	-	-	
<i>Aspergillus niger</i>			Phorbol ester (69.8%) Phytic acid (56.6%)	-	-	
<i>Penicillium miczynskii</i>			Phytic acid (96.5%)	-	-	
<i>Gunninghamella echinulate</i>			Phorbol ester (56.6%) Phytic acid (89.8%)	-	-	
<i>Saccharomyces cerevisiae</i>			Phorbol ester (74.6%) Phytic acid (76.6%)	-	-	
<i>Rhizopus oryzae</i>			Phorbol ester (63.8%) Phytic acid (91.2%)	-	-	
<i>Pleurotus ostreatus</i>	Incubation for 60 days	Phorbol esters	Phorbol ester (62.6%) Phytic acid (73.5%)	-	-	[201]
<i>Candida parapsilosis</i>	Incubated at 30 °C for 24 h	Phorbol esters Curcumin Trypsin inhibitor Saponin	99% 100% 97.8% 64% 94.9%	-	-	[115]

animal feeding trials, with *Candida parapsilosis* showing up to 100% detoxification of phorbol ester in seed cake [115] followed by 99% reduction by *Pleurotus ostreatus* [134]. Among the bacteria studied, *Pseudomonas aeruginosa* showed 100% detoxification of phorbol ester in seed cake [135], compared to lower efficiencies by *Lactobacillus plantarum*, *Propionibacterium* (47.4% reduction) [137] *Bacillus licheniformis* and *Bacillus subtilis* (32.8% reduction) [137]. Although these results suggest strong potential for microbial-detoxification of *J. curcas* seed cake, more information is needed on the life cycle analysis of seed cake and of the environmental impact and economic feasibility for large-scale detoxification.

Genetic engineering approaches have also been explored to generate toxin free seeds of *J. curcas*, and provide a possibility to expand economic feasibility of *J. curcas* seed cake while minimizing the release of toxins into the environment [119,138]. The target of most studies has been the curcins, which are Type I ribosome inactivating proteins and are found in all accession of *J. curcas* including the edible Mexican *J. curcas* varieties which lack phorbol esters [139]. In *J. curcas*, the expression of the *Curcin 1* gene was mainly in endosperm, whereas *Curcin 2A* was expressed in young leaves [139]. Patade et al. [119] conducted RNAi mediated constitutive silencing of *curcin 1* precursor mRNA in *J. curcas* using seed transformation. *Curcin 1* precursor transcript accumulation was compared in mechanically wounded leaves of transgenic and wildtype plants, as mechanical wounding leads to stress-induced expression of the *curcin 1* precursor, which is generally undetectable in leaves under non-stress conditions. The leaves from RNAi plants result showed 98% reduction in the curcin transcript compared to wild type plants. However, it was not reported if there is any reduction in curcin protein content in *J. curcas* seeds from this approach.

The finding that curcin expression increases during biotic and abiotic stresses raises the concern that “curcin-silenced” plants may be more susceptible to stress: Qin et al. [140] showed curcin expression was induced in response to fungal infection (*Pestalotia funerea*, *Curvularia lunata*, *Gibberella zeae*) and abiotic stresses (drought and high temperature), while, ectopic expression of *Curcin 2A* in transgenic tobacco increased disease tolerance against tobacco mosaic virus (TMV) and *Rhizoctonia solani* [141]. A solution to the above issue, by confining the gene silencing to the seeds, was demonstrated by Gu et al. [138]; using a marker-free transgenic *J. curcas* with curcin-deficient seeds this approach provided a solution to curcin toxicity as well as addressing the safety concerns of the marker genes. Gene silencing has also been used to engineer low phorbol ester *J. curcas* by seed specific silencing of two casbene synthase gene (*JcCASA163* and *JcCSD168*), reducing the precursor for phorbol esters [142]. Such gene silenced lines can be used as a parent in *J. curcas* breeding program to generate toxin free varieties. Simultaneous seed-specific silencing of multiple toxic-compound synthetic pathway genes may provide more suitable toxin free varieties.

2.4. Oil quality improvement

Seed oil content is the most economically important attribute of *J. curcas*. Plant oils are mainly composed of triacylglycerols (TAGs), molecules that comprise three fatty acid chains esterified to glycerol. To use as biodiesel, the seed oil is trans-esterified to fatty acid methyl esters [1]. We previously reported the oil content in *J. curcas* seed as around 23.70–46.60% [1]. As in other oleaginous plants, oil biosynthesis in *J. curcas* mainly occurs in the plastid and endoplasmic reticulum [143] (Fig. 3). Inside the plastids, de novo fatty acid synthesis starts with condensation of Acetyl-CoA to Malonyl-ACP, followed by a series of desaturation and hydrolysis reactions [reviewed in 144]. After synthesis, the free fatty acid is exported to the cytoplasm where it is converted into acyl-CoA. The acyl-CoA is transported to the endoplasmic reticulum and is acylated at the sn-1, sn-2, and sn-3 positions of glycerol-3-phosphate to produce TAG via an acyl-CoA dependent pathway known as the Kennedy pathway [145]. An acyl-CoA-independent pathway also exists, where fatty acids are added to

membrane lipids at the plastid envelope and/or in the endoplasmic reticulum via conversion of Diacylglycerol (DAG) to phosphatidylcholine (PC), an interchangeable process catalysed by choline phosphotransferase. PC acts as a donor of fatty acid for DAG to form TAG via Phospholipid: diacylglycerol acyltransferase (PDAT) [146]. After TAG biosynthesis is completed, oil droplets are accumulated as oil bodies (oleosome) in the cytoplasm.

There has been extensive progress in the exploration and characterization of genes associated with oil biosynthesis in *J. curcas*. Examples include: heteromeric subunit genes of ACCase (responsible for carboxylation of acetyl-CoA to produce malonyl-CoA) [147]; stearyl-ACP (involved in the desaturation of stearyl-ACP to Oleoyl-ACP [148]; two beta-ketoacyl-acyl carrier protein synthase genes *KASIII* [149] and *KAS II* [150] (associated with acyl chain elongation); acyl-ACP thioesterase gene (*FATB1*) (involved in the release of fatty acid) [151] and fatty acid desaturase genes *JcFAD2* and *JcFAD3* (involved in desaturation of oleic acid to linoleic acid and α -linolenic acid) [152]. *J. curcas* genes that have been associated with the Kennedy pathway include *GPAT* (Glycerol-3-phosphate acyltransferase) [153], *DGAT1* [154] and *DGAT2* (diglyceride acyltransferase) [155] and *PDAT* [156].

Fuel properties of *J. curcas* biodiesel reported in different geographical location are summarised in Table 3. Some important factors that are restricting the adoption of *J. curcas* biodiesel worldwide are its poor cold-flow properties, high viscosity, high oxidation rate and increased emission of nitrogen oxides (NO_x) [reviewed in 157] relative to conventional fossil fuels. Cold flow properties and viscosity can be improved by decreasing saturated fatty acid levels, whereas, improving oxidative stability and reducing NO_x emission [reviewed in 158] requires a decrease in the proportions of unsaturated and poly-saturated fatty acids in the oil [reviewed in 159]. However, high unsaturation has a negative impact on the ignition quality of fuel [reviewed in 159] making it very challenging to find the ideal fatty acid profile that will deliver biodiesel with optimum parameters. A promising approach is to use oil with a high mono-unsaturated fatty acid content. We observed in our previous study that the fatty acid content in *J. curcas* mainly comprises the monosaturated fatty acid, oleic acid (41.48–46.1%) followed by the polysaturated fatty acid, linoleic acid (30.1–35.2%) [1]. Recent studies have reported increased oil content and modified fatty acid profiles from *J. curcas* based on the genetic manipulation of plants: Ectopic expression of *JcFATB1* (Fatty acyl-acyl carrier protein thioesterase) [151] and *JcKASIII* (β -ketoacyl-acyl carrier protein synthase II) [160] cDNA in Arabidopsis resulted in an increased content of palmitate and reduced levels of unsaturated fatty acids in seed oil. Ectopic expression of *JcDGAT1*-cDNA in Arabidopsis produced an increase in oil content by 30–40% and increase of linolenic acids in the TAG [154], whereas, expression of Arabidopsis *DGAT1* in *J. curcas* was reported to increase oleic acid content in TAG [81]. RNA Silencing approaches to reduce expression of oil biosynthesis genes have also showed some promise: silencing of *FAD2* (fatty acid desaturase 2) in *J. curcas* showed a hike in oleic acid content by more than 78% [161] while silencing of sugar-dependent 1 (*JcSDP1*), (an enzyme responsible for the first step of TAG degradation during seed germination) showed increase in TAG accumulation in *J. curcas* [162]. The availability of a whole genome sequence for *J. curcas* [143,163–166] provide more accessibility to oil biosynthetic genes known from their homologs in other plants, which can accelerate the development of trait-specific markers and the engineering of *J. curcas* varieties with desirable oil quality traits. In particular a clear characterization and evaluation of the functional application of genes of key enzymes involved in TAG synthesis are needed.

2.5. Advances in knowledge of abiotic stress biology of *J. curcas*

J. curcas has been reported to be effected by several types of abiotic stress such as high salinity, low temperature, drought and water logging, however its responses to such stresses are still poorly explored [34]. The impact of drought and salinity on *J. curcas* cultivation must be




Fig. 3. Schematic representation of TAG bio-synthesis ([143]; [226,227]). For simplicity, all possible reactions are not shown. The red ink represents the genes which are isolated and characterized in *J. curcas*. ACCase: acetyl-CoA carboxylase; PEP: Phosphoenolpyruvate; HAR: Hydroxyacyl-ACP dehydratase; KAR: β -ketoacyl-ACP reductase; KAS: β -ketoacyl-ACP synthase; PT: Palmitoyl Thioesterase; ST: Stearoyl Thioesterase; OT: Oleoyl Thioesterase; FAD: Fatty acid desaturases; G3P: Glycerol-3-phosphate; GPAT: Glycerol-3-phosphate acyltransferase; LPA: Lysophosphatidic acid; PA: Lysophosphatidic acid; LPAT: Lysophosphatidic acid acyltransferase; DAG: Diacylglycerol; TAG: Triacylglycerol; PDAT: Phosphatidyl choline transferase; PC: Phosphatidyl choline; CPT: Choline phosphotransferase; LPC: Lysophosphatidyl choline.

known as a basis to standardize irrigation practices for large-scale cultivation. Gao et al. [167] analysed the impact of salinity stress on *J. curcas* seedlings under different NaCl concentrations (50–200 mM) and reported significant reduction in biomass compared to control seedlings. Salinity stress is associated with leaf yellowing, foliar damage and reduction in growth and biomass in *J. curcas*. However, leaf Na^+ (23.6–34.6 g kg DW $^{-1}$) and Cl^- concentrations (11.9–21 g kg DW $^{-1}$) were reported to be much higher than those detected in most glycophytes [168]. On the other hand, water stress conditions (drought) have a negative effect on seed productivity and oil content in field grown *J. curcas* [49]. Extreme drought conditions are reported to initiate leaf drop and limited root growth in *J. curcas* [19]. However, the higher level of proline synthesis recorded for drought-exposed *J. curcas* [127] may also help to explain the short drought-recovery periods (6–7 days) observed in terms of plant growth, stomatal structure and photochemical parameters [169,170]. At the other extreme, water-logging for a duration of 10 days resulted in reduced total biomass, leaf photosynthetic rate and carbohydrate content in both leaves and roots of *J. curcas* [171]. As most studies, have been carried out in controlled environments and with *J. curcas* seedlings, the extrapolation of data for field conditions still remains to be validated. In addition, no systematic

assessment has been done to determine the impact of salinity stress on seed productivity. Therefore, further studies under field conditions and with mature *J. curcas* plants are needed to establish the impacts and to support the development of remedial measures.

At the molecular genetic level of abiotic stress biology, a number of transcription factors from *J. curcas*, have been associated with the regulation of stress-responsive genes [172,173]. The over-expression of *JcDREB*, a putative AP2/EREBP (ethylene responsive element binding protein) domain-containing transcription factor gene isolated from *J. curcas*, resulted in enhanced salt and freezing-stress tolerance in the model plant *Arabidopsis* [172]. Transgenic studies in *J. curcas* have also shown promise for salt and drought-stress tolerance [83,174]. The constitutive expression of a vacuolar Na^+/H^+ antiporter gene from the halophyte plant *Salicornia brachiata*, in *J. curcas* provided increased salt tolerance (at 200 mM NaCl exposure) compared to wild type plants [83]. Genome-wide profiling of MYB family transcription factors in *J. curcas* led to the identification of 128 MYB genes, among which four subfamilies of domain repeats were observed including, 123 R2R3-MYB, four R1R2R3-MYBs and one 4R-MYB [173]. From these, over-expression of the nuclear localized *JcMYB2* in *Arabidopsis*, improved tolerance to salt and cold stress [173]. Tissue specific expression

Table 3
Properties of biodiesel from *J. curcas* from different geographical locations.

Geographical location	Oil content (%)	Viscosity (cst)	Density (g/cm 3)	Cloud point (°C)	Pour point (°C)	Flash point (°C)	Calorific value (MJ/kg)	References
Nicaragua (South America)	–	4.84	0.88	–	–	191	–	[202]
Faisalabad (Pakistan)	30–40	4.80 \pm 0.17	0.88	10 \pm 0.1	6 \pm 0.2	188 \pm 3.0	–	[203]
Sichuan (South west china)	–	4.06–5.13	0.82	–	–	164–166	–	[204]
Nigeria (Africa)	47.25	–	0.91	–	–	–	–	[205]
Malaysia	60	2.83	0.91	–	–	190.5	–	[206]
Nigeria	–	9.60	0.88	–	–	200	48.31	[207]
Brazil	31.6	4.02	0.88	–	– 5	117	40.31	[208]
China	–	3.6	–	1	–	153	–	[209]
Yunnan (South west China)	–	3.89	0.88	–	– 5	186	41.72	[210]
Nigeria	–	4.73	0.86	–	–	184	–	[211]
Assam (India)	23.70–46.60	6.10–6.70	0.83–0.85	8	2	185	39.56–41.09	[1]
Malaysia	–	4.7227	0.86	5	3	182.5	39.827	[157]
Honduras (Central America)	–	2.7–3.0	–	3.8–4.3	–	–	–	[212]
Udaipur (India)	–	5.48 \pm 0.06	0.86	–	–	–	39.47	[213]

profiles of *JcMYB2* in *J. curcas* showed it to be ubiquitously expressed throughout the plant, with higher transcript accumulation in roots [173]. Genome-wide analyses of salt [175], drought [176] and cold [177]-responsive transcriptomes of *J. curcas*, have identified numerous stress-responsive genes with potential for application in developing abiotic-stress tolerant varieties of the crop.

J. curcas is susceptible to water stress with poor performance under drought or water logging and research on transcription factors has also advanced knowledge in this area. *J. curcas* lines over-expressing Arabidopsis B subunit of the nuclear factor Y (*AtNF-YB1*) gene, showed enhanced content of the osmoprotectant, glycine betaine [174], although the impact of this on stress-tolerance was not studied. The adjustment of *J. curcas* roots against waterlogging-stress, explored at the transcriptome level via high-throughput RNA-sequencing (RNA-seq), identified the involvement of transcription factor families including AP2/ERF, MYB, and WRKY [178]. From differential gene expression data, it could be shown that water logging triggers responses in genes associated with hypoxia, anaerobic respiration, deterioration of cell wall biogenesis, carbohydrate production and growth in *J. curcas* [178]. Since adverse climatic conditions are not only the norm for non-agricultural soils, but are expected to extend their range in the light of climate change, the findings on abiotic stress responses in *J. curcas* can provide new scopes for developing tolerant varieties, but also for other crops in the future.

2.6. Impact of biotic stress and measures to enhance cultivation of *J. curcas*

J. curcas seed oil contains compounds that are toxic to many microorganisms, insects and pests [179]. Despite the toxicity of the seeds, *J. curcas* plants are still susceptible to several pests and diseases, as a result of which, various measures have been explored for pest and disease control, some examples are shown in Table 4. The insect pest susceptibility of a plant species highly depends on the geographic location and environmental conditions, as these influence the presence of the pest or pathogen and the success of its infestation or infection. The major pests affecting *J. curcas* cultivars belong to the orders Lepidoptera, Hemiptera and Heteroptera. These pests have been reported to cause leaf mining, premature abortion of inflorescence and fruits and to account for 60–80% damage in large-scale plantation (Table 4). Spraying of insecticides can address these issues, however, may cause undesirable environmental issues including ground-water contamination and air pollution. The residues of chemical insecticides are toxic to humans and domestic animals [180] as well as to non-target and beneficial insects like honeybees, which play a very important role in *J. curcas* pollination and hence effect seed productivity [181]. A further complication is that poor management of insecticide application can result in the emergence of resistant insect biotypes [182].

Eco-friendly alternatives for pest and disease control have been explored, including use of biopesticides and other biocontrol measures, as these are biodegradable and have a reduced impact on the environment. Djimmy et al. [183] recently suggested controlling insect pests of *J. curcas* by exploring natural enemies of the insect: african shield bug (*Calidea panaethiopica*), a polyphagous heteroptera of the Scutelleridae family, feeds on *J. curcas* flowers, fruit, and seeds thereby causing huge loss in seed production in the South-Sudan Zone of Burkina Faso. Three parasites (*Trissolcus basalis*, *Psixstriaticeps* and *Gryon* sp.) were identified to infect the eggs of *C. panaethiopica* [183] with a peak of 43% parasitism during June to September. However, to exploit natural enemies associated with this insect pest, the biology, ecology, life cycle and host range of the predators should be thoroughly studied as well. Another viable alternative to chemical insecticides is the use of microbes e.g. *Bacillus thuringiensis* (*Bt*) [184]. Different strains of *Bt* are effective against specific Lepidopteran, Dipteran, Coleopteran, Hymenoptera, Homoptera, Orthoptera and Mallophaga insects as well as against nematodes and mites, which has led to their extensive use in crop protection [reviewed in 185]. However, two major limitations for

using *Bt* spore formulations in crop protection are the limited use in wet climates or during rainy seasons, as formulations are easily washed away and the poor effectivity of sprayed spores against insect pests residing within plant tissues [186]. Transgenic crops expressing bacterial *Bt* toxins, have successfully addressed these challenges and are some of the most widely deployed and commercially successful plant biotech products. We previously reported transgenic *J. curcas* expressing the *Bt* toxin Cry1Ac protein which is toxic to the lepidopteran insect *Spodoptera litura* [187]. A marker-free transgenic line of *J. curcas* expressing hybrid *Bt* δ -endotoxin protein Cry1Ab/1Ac showed promising toxicity (80–100% mortality rate) towards the lepidopteran insect *Archips micaceanus* [188]. While these studies in *J. curcas* are at an early stage, the overwhelming success of the BT trait in other crops encourages us to believe that insect-resistant *J. curcas* lines are feasible and can increase seed yield in large-scale plantation while reducing the cost and environmentally detrimental effects of high pesticide use.

Fungal pathogens also can reduce *J. curcas* yields. While *J. curcas* leaf extract has shown inhibitory effect on mycelial growth of *Usarium graminearum*, *Pyricularia oryzae*, *Phytophthora nicotianae* and *Phytophthora capsica* [189], *J. curcas* has been reported to be susceptible to several other fungal parasites, including *Fusarium oxysporum* (wilting), *Rhizoctonia bataticola* (root rot), *Alternaria alternat* (leaf spot), *Fusarium moniliforme* (root rot) and *Colletotrichum gloeosporioides* (Anthracnose) (Table 4). Spraying of fungicides such as carbendazim provided 100% inhibition of *Fusarium oxysporum* [190] and a combination of Bavistin and Vitavax at 50 ppm completely inhibited mycelial growth of *Rhizoctonia bataticola* (*R. bataticola*) [191]. However, as with insecticides, spraying of fungicides possesses environmental and health hazards, spurring research into alternate solutions. Kumar et al. [191] reported an integrated approach using plant extracts and pathogenic fungi together to reduce use of chemical fungicides: A combination of Bavistin (2 g/kg seed) with *Azadirachta indica* (neem extract) (20%) resulted in 67.3% inhibition of *R. bataticola* growth, while, a combination of *Trichoderma harzianum* (*T. harzianum*) (15 g mycelial mat /kg seed) and Bavistin showed 54.2% reduction in *R. bataticola* growth [191].

Other than insect and fungal parasites, *J. curcas* is also vulnerable to viruses, including various strains of *J. curcas* mosaic virus and Indian cassava mosaic virus (Table 4). There are several reports of virus-resistant plant lines which express viral coat protein or hairpin dsRNA with sequences homologous to viral genes [192]. Ye et al. [83], first reported development of transgenic *J. curcas* plants resistant to the geminivirus, Indian cassava mosaic virus (ICMV) via expression of a hairpin dsRNA of five ICMV DNA-A genes. Although initially laborious and expensive to generate, resistant transgenic lines can be used in as a parent in *J. curcas* breeding programs to rapidly multiply useful resistance traits. An alternate rapid development approach is to directly treat plants with dsRNA, which can be inexpensively made as a crude extract from bacteria expressing dsRNA with sequences homologous to viral mRNA [193,194]. The spraying should be done before the expected outbreaks of virus infection [194], such as when a vector insect infestation is first detected. This approach greatly depends on weather conditions, however, recent research on delivery of dsRNA loaded in clay nanoparticles (BioClay) showed the dsRNA to be stable and effective against crop viruses for an extended period of 20 days [195]. Thus, this approach could also be deployed for *J. curcas* in the future. The emerging research on biological approaches to *J. curcas* improvement are encouraging and attractive, as biological approaches are generally environmental safe. However, further research related to the specific pests and diseases of *J. curcas* is needed to develop solutions that are suitable for large scale deployment and are economically viable and stable.

3. Conclusions and recommendations

J. curcas has been shown to be a renewable source for sustainable

Table 4
Summary of biotic stress, area of occurrence and measures taken to control damage/infection in *J. curcas*.

Causal organism	Order	Damage/Symptom	Control suggested	Place reported	Reference
Insect					
<i>Spodoptera litura</i>	Lepidoptera	Devour the leaves, effect overall growth of the plant.	-	India (Madhya Pradesh)	[214]
<i>Scutellera perplexa</i>	Hemiptera	Fruit feeding led to premature fruit and seed abortion and reduction in yield.	-	India (Lucknow)	[18]
<i>Maonellucoccus hirsutus</i>	Hemiptera	Curling and contortion of leaves, stunted plant, deformed fruit, high abundance of insects can cause death of the tree.	-	India (Lucknow)	[215]
<i>Stomphastis thraustica</i>	Lepidoptera	Insect larvae feed on leaf blades by digging tunnels into them, leading to large brown patches on the leaves.	-	India (Secunderabad, Jhansi)	[216,217]
<i>Pempelia morosalis</i>	Lepidoptera	Fed on leaf and pollen causing damage to the inflorescences, stems and fruit.	-	Malaysia	[218]
<i>Calidea panaethiopica</i>	Heteroptera	Fed on the flowers and fruits of <i>J. curcas</i> . Attacked flowers turned dry and attacked fruits often have malformed or empty seeds.	-	China (Panzhihua)	[219]
				Belgium (Lower Valley of the Senegal River)	[220]
				Belgium (Lower Valley of the Senegal River)	[220]
				Belgium (Lower Valley of the Senegal River)	[220]
				West Niger	[221]
				Burkina Faso	[183]
Fungi					
<i>Fusarium oxysporum</i>	Hypocreales	Wilting, chlorosis, necrosis, premature leaf drop, browning of the vascular system and stunted plants.	Spraying of carbendazim (200 µg/mL) showed 98% growth inhibition.	China (Hainan)	[190]
<i>Rhizoctonia bataticola</i>	Cantharellales	Root rot, symptoms of yellowing and dropping of leaves, dark brown lesions at collar region which spreads downwards in the plant.	Fungicides such as Bavistin and Vitavax at 50 ppm showed 100% inhibition of mycelial growth. Bavistin (2 g/kg seed) + neem extract (20%) showed 67.3% reduction, <i>Trichoderma harzianum</i> (1.5 g per seed) + Bavistin and neem extract + T. harzianum showed 44.0% reduction	India (Haryana)	[191]
<i>Alternaria alternata</i>	Pleosporales	Amphigenous, irregular infection spots (1–2 cm) appear on both surfaces of leaf, which turns dark in the later stage.	-	India (Uttar Pradesh)	[222]
<i>Fusarium moniliforme</i>	Hypocreales	Yellowing of leaves leading to defoliation, drying of plants from tip downwards and ultimately death of the plant. On up- rooting the affected plants, browning/blackening of the roots was observed.	-	India	[223]
<i>Colletotrichum gloeosporioides</i>	Ascomycetes	Dark, water soaked lesions on stems, leaves or fruit.	-	Brazil and Cape Verde	[224]
Virus					
Mosaic virus	Begomovirus	Mild to severe mosaic, marked reduction in leaf size, rolling of leaf margins and puckering of the leaf surface. Chlorotic areas of irregular shapes were present between the secondary veins.	-	India	[225]
Jatropha mosaic virus	Geminivirus	Mild to severe mosaic, marked reduction in leaf size, rolling of leaf margins.	Transgenic <i>J. curcas</i> plants with resistance to ICMV via expression of a hairpin dsRNA with sequences homologous to five key ICMV DNA-A genes showed resistance.	Singapore	[83]
Indian cassava mosaic virus	Begomovirus	Yellow-green mosaic, curling, malformation and size reduction. In addition, infected plants also displayed other symptoms including shortened internodes, stunting of plant stature, fewer flowers which were partially or completely sterile.	-	India (Dharwad)	[10]

energy production, with good potential to address issues related to economic growth. Many of the valid concerns over various shortcomings of the crop arise from the requirement for production using marginal and contaminated soils so as to avoid competition for land for agricultural crop production. Despite of many challenges, the potential for *J. curcas* as biodiesel plant remains promising in the light of new knowledge and biotechnology. This review highlights the advancements from biological studies of *J. curcas* that are pertinent to application of this crop as a future sustainable biodiesel source. A summary of the main findings, from which our recommendations for further research have been made are as follows:

3.1. Improvement of seed yield of *J. curcas* via soil enrichment

Growing non-edible biodiesel crops in low nutrient and marginal soils offers a sustainable alternative approach to reclaim those lands while avoiding competition with food crops for land use. However, for cost-effective and profitable plantation, soil enrichments procedures are required. Although chemical fertilizers promote plant growth, their application leads to groundwater contamination, greenhouse gas emissions and disturbs the microflora and fauna of the soil. Successful alternative approaches are available via use of beneficial microbes such as AMF, endophytes and PGPR as biofertilizers. Application of AMF, endophytes and PGPRs led to enhanced seed production in *J. curcas* in low nutrient and stress conditions. However, the rhizosphere environment differs from that of internal plant tissues and also in different soil conditions. For example, the variations in light intensity, soil temperature, pH, nutrient availability, and the interaction with other organisms in the rhizosphere may influence performance. Therefore, it is recommended that indigenous species of AMF and PGPR are used in each locality. There are still several challenges such as optimising systems for low-cost mass production, and improving the viability of microbial inoculant during storage. In addition, use of elite cultivars of *J. curcas* with improved stress-tolerance will ameliorate some of the yield-reduction that results from poor nutrition and contaminants in marginal soils.

3.2. Manipulation of inflorescence development to enhance seed yield

Plants producing inflorescences with an increased female flower ratio is a highly desirable trait for biodiesel crops. While exogenous application of plant growth regulators such as gibberellins, cytokinin and thidiazuron have shown substantial increase in female flowers, the process is labour intensive and success is weather-dependent. Recent advancement in genomics and transcriptomics, provides more options for improvement in female flower numbers in *J. curcas* via manipulation of plant growth regulator biosynthesis pathways or key sex-related genes responsible for sex differentiation at the developmental biology level. The over expression of flowering locus T homolog (*JcFT*) [97] and *LEAFY* (*JcLFY*) [98] showed promise in this respect and produced plants with higher numbers of female flowers. Average seed yield in *J. curcas* can be improved by providing suitable pollination services to optimise fruit set via cross pollination using honey bee species that have been recorded as the most efficient pollinators. Therefore, the strategic establishment of native bee hubs inside plantation areas is recommended.

3.3. Detoxification of seed cake to aid economic value of *J. curcas* plantations

Seed cake detoxification using fungal and bacterial strain has shown promising results. The fungi *Candida parapsilosis* [115] and *Pleurotus ostreatus* [134] and the bacteria *Pseudomonas aeruginosa* [135] were observed as the most efficient bioagents for detoxification. Following detoxification, seed cakes can be used as a component of biofertilizers and animal feed. However, for profitable industrial implementation of

seed cakes life cycle analysis, environmental impact assessment and cost analysis are needed. In addition, the toxin-free seeds from genetically modified *J. curcas* lines can aid in economic feasibility by reducing the processing costs for toxin-free seed cakes for fertilizers and animal feed.

3.4. Modification of fatty acid composition for improvement of biodiesel properties

Oil high in mono-saturated fatty acid and low in both saturated and polysaturated fatty acid is always desirable as biodiesel feedstock to address issues such as cold flow properties, oxidative stability and NOx emissions. Some progress has been achieved through the manipulation of fatty acid biosynthesis genes, such as reduced poly-unsaturated fatty acid levels in model plants through the over-expression of fatty acyl-carrier protein thioesterase (*JcFATB1*) [151] and β -ketoacyl-acyl carrier protein synthases II (*JcKASIII*) [160]. Increased monosaturated fatty acid (oleic acid, 18:1) content and the reduction in both saturated and polysaturated fatty acid in *Jatropha* seed oil were also demonstrated after over-expression of *Arabidopsis* diglyceride acyltransferase *DGAT1* [81] and silencing of fatty acid desaturase 2 (*FAD2*) [157] in *J. curcas*. Further improvement in cold flow properties could be achieved by increasing the short chain mono-saturated (palmitoleic acid, 16:1) and saturated fatty acid content of *J. curcas* seed oil via silencing fatty acid desaturases responsible for conversion of long chain saturated (e.g. 18:2, 18:3) or mon-unsaturated (e.g. 18:1, 20:1) fatty acid. To date, relatively few genes have been studied and further functional characterization, in particular for TAG biosynthesis genes, can advance this approach to *Jatropha* improvement. Recent publication of the whole genome sequence for *J. curcas* can facilitate both genetic engineering and molecular breeding approaches to develop *J. curcas* lines with desired oil quality traits.

3.5. Improving abiotic stress tolerance

Abiotic stress factors such as salt, drought and waterlogging limit seed productivity of *J. curcas*. The early approach to combat abiotic stress was to develop stress-tolerant cultivars using conventional breeding. However, no modified stress-tolerant cultivars have been reported via breeding as the process is not only labour intensive but also time consuming for a perennial crop such as *J. curcas*. Early research using a genetic modification approach to overexpress a vacuolar Na⁺/H⁺ antiporter from *Salicornia brachiata* in *J. curcas* showed promise, including a *J. curcas* variety with increased tolerance to salt stress, however field validation of novel hybrids will be needed before commercial viability is assured. Meanwhile, stress-responsive gene discovery is providing several promising target genes to test (e.g. the transcription factors *JcDREB*, *JcMYB*, *AP2/ERF* and *WRKY*) and it is anticipated that, as for other crops, new varieties that can produce good yields on marginalised soils will emerge in time, if given the needed research support for development and evaluation.

3.6. Improving biotic stress tolerance

Despite the toxins found in the seed, *Jatropha* is susceptible to numerous pests and diseases. Two potentially complementary biological approaches have promise here, which are biopesticide/biological control of pests, and genetic improvement of resistance traits in the plant, leveraging new knowledge on pest and disease responsive genes in *J. curcas* as well as from other crops. Another potential approach is to exploit natural parasitic relationships among insect pests, which has been reported for the heteropteran insect *Calidea panaethiopica*. Knowledge on biology, ecology, life cycle and host range of the predators should be obtained prior to application. Development of insect resistance is another stable approach to control insect manifestation, by deploying suitable candidate genes. Other biotic stresses such as fungal

attack can be diminished by application of combination of pathogenic fungus and plant extracts. Viral attacks can be suppressed by spraying dsRNA loaded in bio-clay containing homologous sequences of viral mRNA or developing virus resistance transgenic *J. curcas*. However, both gene silencing approaches have pros and cons, direct treatment with dsRNA is rapid but able to give protection only up to 20 days after each spraying, while the development of transgenic resistant plant lines takes longer but can provide more stable protection of the plant throughout the lifetime.

4. Future prospects

Based on the progress in biological research together with newly emerging technology, there are excellent prospects to advance *J. curcas* as a bioenergy crop for use on marginal soils. Firstly, the use of bio-agents including microbes as biofertilizers and biopesticides has shown promising results for enriching soil fertility and in providing protection against biotic and abiotic stresses, with the additional benefits of reduced input cost, improved environmental safety and target-specificity. The exploration of the molecular basis of host interactions with beneficial microbes will further facilitate establishment of a stable eco-friendly and sustainable *J. curcas* plantation, avoiding or at least reducing the use of chemical fertilizers, insecticides or pesticides.

Secondly, the availability of the whole genome sequence, efficient transformation systems and genetically modified lines of *J. curcas*, offer exciting opportunities to develop elite commercial cultivars with high yield along with tolerance to biotic and abiotic stresses. Early studies have shown that both over-expression and gene silencing methods can address the various limitations of this crop. However, since the deployment of genetically modified crops, still faces issues of public acceptance in several countries, alternative approaches can be explored such as direct application of dsRNA or gene editing with site specific nucleases [196]. The emergence of highly precise genome editing techniques which involve specific alterations of single or multiple genetic loci, has been effective as a novel targeted approach to crop mutagenesis which in some legislations is not regulated as a genetic modification [reviewed in 197]. It is attractive to think that site-specific nucleases such as clustered regularly interspaced short palindromic repeat (CRISPR)-associated endonuclease 9 (CRISPR/cas9) that have shown successful results in plant improvement [198] can also be used in *J. curcas*. While the efficacy of these approaches in genetic improvement of *J. curcas* has not been demonstrated so far, it is feasible that site-directed mutagenesis of precursor genes of curcin and phorbol ester could generate varieties of *J. curcas* with toxin-free seed. Similarly, gene silencing and editing can be used to develop elite cultivars with many of the currently lacking traits, and also provide the possibility of combining abiotic and biotic stress resistance, improved oil quality for biodiesel use and toxin-free seed cake as a commercially valuable by-product. A combinatorial approach using elite lines from conventional breeding programs, complemented by and crossed with those produced via biotechnological tools can expedite crop improvement. Taken together biological information and biotechnological approaches can help to bridge the gaps in productivity for *J. curcas*.

Acknowledgements

Author Purabi Mazumdar acknowledge Prof. Lingaraj Sahoo, Centre for Energy, Indian Institute of Technology Guwahati for introducing her on *Jatropha* related research during her PhD work. (PM, PS and JAH are partially supported by CEBAR Research University grant (RU006–2017 and RU015–2016).

References

- [1] Mazumdar P, Borugadda VB, Goud VV, Sahoo L. Physico-chemical characteristics of *Jatropha curcas* L. of North East India for exploration of biodiesel. Biomass-

- Bioenergy 2012;46:546–54.
- [2] Prasad DR, Izam A, Khan MMR. *Jatropha curcas*: plant of medical benefits. J Med Plants Res 2012;6:2691–9.
- [3] Warra AA. Cosmetic potentials of physic nut (*Jatropha curcas* Linn.) seed oil: a review. Am J Sci Ind Res 2012;3:358–66.
- [4] Openshaw K. A review of *Jatropha curcas* L.: an oil plant of unfulfilled promise. Biomass- Bioenergy 2000;19:1–15.
- [5] Banapurmath NR, Tewari PG, Hosmath RS. Performance and emission characteristics of a DI compression ignition engine operated on Honge, *Jatropha* and sesame oil methyl esters. Renew Energy 2008;33:1982–8.
- [6] Nahar K, Ozores-Hampton M. *Jatropha*: an alternative substitute to fossil fuel. IFAS Publication Number HS1193. Gainesville: University of Florida, Institute of Food and Agricultural Sciences; 2011.
- [7] Divakara BN, Upadhyaya HD, Wani SP, Gowda CL. Biology and genetic improvement of *Jatropha curcas* L.: a review. Appl Energy 2010;87:732–42.
- [8] Wani SP, Chander G, Sahrawat KL, Rao CS, Raghvendra G, Susanna P, Pavani M. Carbon sequestration and land rehabilitation through *Jatropha curcas* (L.) plantation in degraded lands. Agric Ecosyst Environ 2012;161:112–20.
- [9] Chang FC, Ko CH, Tsai MJ, Wang YN, Chung CY. Phytoremediation of heavy metal contaminated soil by *Jatropha curcas*. Ecotoxicology 2014;23:1969–78.
- [10] Gao S, Qu J, Chua NH, Ye J. A new strain of Indian cassava mosaic virus causes a mosaic disease in the biodiesel crop *Jatropha curcas*. Arch Virol 2010;155:607–12.
- [11] Luoma JR. Hailed as a miracle biofuel, *Jatropha* falls short of hype. Yale Environment; 2009. p. 360.
- [12] Ariza-Montobbio P, Lele S, Kallis G, Martinez-Alier J. The political ecology of *Jatropha* plantations for biodiesel in Tamil Nadu, India. J Peasant Stud 2010;37:875–97.
- [13] Kant P, Wu S. The extraordinary collapse of *Jatropha* as a global biofuel. Environ Sci Technol 2011;45:7114–5.
- [14] Singh K, Singh B, Verma SK, Patra DD. *Jatropha curcas*: a ten-year story from hope to despair. Renew Sustain Energy Rev 2014;35:356–60.
- [15] Sop TK, Kagambega FW, Bellefontaine R, Schmiedel U, Thiombiano A. Effects of organic amendment on early growth performance of *Jatropha curcas* L. on a severely degraded site in the Sub-Sahel of Burkina Faso. Agrofor Syst 2012;86:387–99.
- [16] Gangwar M, Sood H, Chauhan RS. Genomics and relative expression analysis identifies key genes associated with high female to male flower ratio in *Jatropha curcas* L. Mol Biol Rep 2016;43:305–32.
- [17] Xiao J, Zhang H, Niu L, Wang X, Lu X. Evaluation of detoxification methods on toxic and antinutritional composition and nutritional quality of proteins in *Jatropha curcas* meal. J Agric Food Chem 2011;59:4040–4.
- [18] Sahai K, Srivastava V, Rawat KK. Impact assessment of fruit predation by *Scutellera perplexa* Westwood on the reproductive allocation of *Jatropha*. Biomass- Bioenergy 2011;35:4684–9.
- [19] Krishnamurthy L, Zaman-Allah M, Marimuthu S, Wani SP, Rao AK. Root growth in *Jatropha* and its implications for drought adaptation. Biomass- Bioenergy 2012;39:247–52.
- [20] Kumar A, Sharma S. An evaluation of multipurpose oil seed crop for industrial uses (*Jatropha curcas* L.): a review. Ind Crops Prod 2008;28:1–10.
- [21] Devappa RK, Makkar HP, Becker K. Nutritional, biochemical, and pharmaceutical potential of proteins and peptides from *Jatropha*: review. J Agric Food Chem 2010;58:6543–55.
- [22] Abhilash PC, Srivastava P, Jamil S, Singh N. Revisited *Jatropha curcas* as an oil plant of multiple benefits: critical research needs and prospects for the future. Environ Sci Pollut R 2011;18:127–31.
- [23] Kamal S, Mammoohan S, Birendra S. A review on chemical and medicobiological applications of *Jatropha curcas*. I Res. J Pharm 2011;2:61–6.
- [24] Nithiyanantham S, Siddhuraju P, Francis G. Potential of *Jatropha curcas* as a bio-fuel, animal feed and health products. J Am Oil Chem Soc 2012;89:961–72.
- [25] Sabandar CW, Ahmat N, Jaafar FM, Sahidin I. Medicinal property, phytochemistry and pharmacology of several *Jatropha* species (Euphorbiaceae): a review. Phytochem 2013;85:7–29.
- [26] Abdelgadir HA, Van Staden J. Ethnobotany, ethnopharmacology and toxicity of *Jatropha curcas* L. (Euphorbiaceae): a review. S Afr J Bot 2013;88:204–18.
- [27] Jain S, Sharma MP. Prospects of biodiesel from *Jatropha* in India: a review. Renew Sustain Energy Rev 2010;14:763–71.
- [28] Parawira W. Biodiesel production from *Jatropha curcas*: a review. Sci Res Essays 2010;5:1796–808.
- [29] Koh MY, Ghazi TIM. A review of biodiesel production from *Jatropha curcas* L. oil. Renew Sustain Energy Rev 2011;15:2240–51.
- [30] Silitonga AS, Atabani AE, Mahlia TMI, Masjuki HH, Badruddin IA, Mekhilef S. A review on prospect of *Jatropha curcas* for biodiesel in Indonesia. Renew Sustain Energy Rev 2011;15:3733–56.
- [31] Ong HC, Mahlia TMI, Masjuki HH, Norhasyima RS. Comparison of palm oil, *Jatropha curcas* and *Calophyllum inophyllum* for biodiesel: a review. Renew Sustain Energy Rev 2011;15:3501–15.
- [32] Abdulla R, Chan ES, Ravindra P. Biodiesel production from *Jatropha curcas*: a critical review. Crit Rev Biotechnol 2011;31:53–64.
- [33] Mofijur M, Masjuki HH, Kalam MA, Hazrat MA, Liaquat AM, Shahabuddin M, Varman M. Prospects of biodiesel from *Jatropha* in Malaysia. Renew Sustain Energy Rev 2012;16:5007–20.
- [34] Li Z, Lin BL, Zhao X, Sagisaka M, Shibasaki R. System approach for evaluating the potential yield and plantation of *Jatropha curcas* L. on a global scale. Environ Sci Technol Lett 2010;44:204–2209.
- [35] Axelsson L, Franzén M, Ostwald M, Berndes G, Lakshmi G, Ravindranath NH. *Jatropha* cultivation in southern India: assessing farmers' experiences. Biofuels

- Bioprod Biorefin 2012;6(3):246–56.
- [36] van Eijck JAJ, Smeets E, Romijn HA, Balkema AJ, Jongschaap REE. *Jatropha* assessment. Agronomy, socio-economic issues, and ecology. Facts from literature. NL Agency; 2010.
- [37] Jingura RM. Socio-economy, agro-ecological zones, agronomic practices and farming system of *Jatropha curcas* L. in Sub-Saharan Africa. In *Jatropha*. Challenges for a New Energy Crop. New York: Springer; 2012. p. 53–69.
- [38] van Eijck J, Romijn H, Balkema A, Faaij A. Global experience with *Jatropha* cultivation for bioenergy: an assessment of socio-economic and environmental aspects. *Renew Sustain Energy Rev* 2014;32:869–89.
- [39] Kumar S, Chaube A, Jain SK. Sustainability issues for promotion of *Jatropha* biodiesel in Indian scenario: a review. *Renew Sustain Energy Rev* 2012;16:1089–98.
- [40] Achten WM, Nielsen LR, Aerts R, Lengkeek AG, Kjaer ED, Trabucco A, Hansen JK, Maes WH, Graudal L, Akinnifesi FK, Muys B. Towards domestication of *Jatropha curcas*. *Biofuels* 2010;1:91–107.
- [41] Johnson TS, Eswaran N, Sujatha M. Molecular approaches to improvement of *Jatropha curcas* Linn. as a sustainable energy crop. *Plant Cell Rep* 2011;30:1573–91.
- [42] Moniruzzaman M, Yaakob Z, Khatun R. Biotechnology for *Jatropha* improvement: a worthy exploration. *Renew Sustain Energy Rev* 2016;54:1262–77.
- [43] Mukherjee P, Varshney A, Johnson TS, Jha TB. *Jatropha curcas*: a review on biotechnological status and challenges. *Plant Biotechnol Rep* 2011;5:197–215.
- [44] Caesar SA, Ignacimuthu S. Applications of biotechnology and biochemical engineering for the improvement of *Jatropha* and Biodiesel: a review. *Renew Sustain Energy Rev* 2011;15:5176–85.
- [45] Wakandigara A, Nhamo LRM, Kugara J. Chemistry of phorbol ester toxicity in *Jatropha curcas* seed-A review. *Int J Biochem Res Rev* 2013;3:146.
- [46] Baldini M, Raranciu S, Vischi M. Toxicity of *Jatropha curcas* L. as a biofuel crop and options for management plant. *Sci Rev* 2013:193.
- [47] Maghuly F, Laimer M. *Jatropha curcas*, a biofuel crop: functional genomics for understanding metabolic pathways and genetic improvement. *Biotechnol J* 2013;8:1172–82.
- [48] Achten WM, Sharma N, Muys B, Mathijs E, Vantomme P. Opportunities and constraints of promoting new tree crops—lessons learned from *Jatropha*. *Sustainability* 2014;6:3213–31.
- [49] Edrisi SA, Dubey RK, Tripathi V, Bakshi M, Srivastava P, Jamil S, Singh HB, Singh N, Abhilash PC. *Jatropha curcas* L.: a crucified plant waiting for resurgence. *Renew Sustain Energy Rev* 2015;41:855–62.
- [50] Yong JWH, Ng YF, Tan SN, Chew AYL. Effect of fertilizer application on photosynthesis and oil yield of *Jatropha curcas* L. *Photosynthetica* 2010;48:208–18.
- [51] Melillo JM, Reilly JM, Kicklighter DW, Gurgel AC, Cronin TW, Paltsev S, Felzer BS, Wang X, Sokolov AP, Schlosser CA. Indirect emissions from biofuels: how important? *Science* 2009;326:1397–9.
- [52] Achten WMJ, Verchot L, Franken YJ, Mathijs E, Singh VP, Aerts R, Muys B. *Jatropha* bio-diesel production and use. *Biomass- Bioenerg* 2008;32:1063–84.
- [53] Nordborg M, Davis J, Cederberg C, Woodhouse A. Freshwater ecotoxicity impacts from pesticide use in animal and vegetable foods produced in Sweden. *Sci Total Environ* 2017;581–582:448–59.
- [54] Bücking H, Kafle A. Role of arbuscular mycorrhizal fungi in the nitrogen uptake of plants: current knowledge and research gaps. *Agronomy* 2015;5:587–612.
- [55] Smith SE, Facelli E, Pope S, Smith FA. Plant performance in stressful environments: interpreting new and established knowledge of the roles of arbuscular mycorrhizas. *Plant Soil* 2010;326:3–20.
- [56] Charoenpakdee S, Phosri C, Dell B, Lumyong S. The mycorrhizal status of indigenous arbuscular mycorrhizal fungi of physic nut (*Jatropha curcas*) in Thailand. *Mycosphere* 2010;1:167–81.
- [57] Kumar A, Sharma S, Mishra S. Influence of arbuscular mycorrhizal (AM) fungi and salinity on seedling growth, solute accumulation, and mycorrhizal dependency of *Jatropha curcas* L. *J Plant Growth Regul* 2010;29:97–306.
- [58] Kamalvanshi M, Kumar A, Jha A, Dhyani SK. Occurrence of arbuscular mycorrhizal fungi in rhizosphere of *Jatropha curcas* L. in arid and semi arid regions of India. *J Microbiol* 2012;52:492–4.
- [59] Behera SK, Srivastava P, Tripathi R, Singh JP, Singh N. Evaluation of plant performance of *Jatropha curcas* L. under different agro-practices for optimizing biomass—a case study. *Biomass- Bioenerg* 2010;34:30–41.
- [60] Moreira BC, Rodrigues AL, Oliveira SF, Miguel PSB, Bazzolli DMS, Sturmer SL, Kasuya MCM. Arbuscular mycorrhizal fungi in the *Jatropha curcas* rhizosphere. *Afr J Microbiol Res* 2015;9:1060–74.
- [61] Leye EH, Ndiaye M, Ndiaye F, Diallo B, Sarr AS, Diouf M, Diop T. Effet de la mycorrhization sur la croissance et le développement de *Jatropha curcas* L. *Rev Des énergies renouvelables* 2009;12:269–78.
- [62] Aggangan NS, Albano AB, Kasahara ES, Ragragio EM. Survival, growth and Cu accumulation by non-mycorrhizal and mycorrhizal *Jatropha curcas* L. seedlings or cuttings in a grassland and in mine tailing soils. *J Environ Sci Manag* 2013;16:63–76.
- [63] Aggangan N, Cadiz N, Llamado A, Raymundo A. *Jatropha curcas* for Bioenergy and Bioremediation in Mine Tailing Area in Mogpog, Marinduque, Philippines. *Energy Procedia* 2017;110:471–8.
- [64] Castillo CG, Borie F, Oehl F, Sieverding E. Arbuscular mycorrhizal fungi biodiversity: prospecting in Southern-Central zone of Chile. A review. *J Soil Sci Plant Nutr* 2016;16:400–22.
- [65] Jha CK, Saraf M. Invitro evaluation of indigenous plant growth promoting rhizobacteria isolated from *Jatropha curcas* rhizosphere. *Int J Genet Eng Biotechnol* 2011;2:91–100.
- [66] Madhaiyan M, Peng N, Te NS, Hsin C, Lin C, Lin F, Reddy C, Yan H, Ji L. Improvement of plant growth and seed yield in *Jatropha curcas* by a novel nitrogen-fixing root associated Enterobacter species. *Biotechnol Biofuels* 2013;6:140.
- [67] Qin S, Miao Q, Feng WW, Wang Y, Zhu X, Xing K, Jiang JH. Biodiversity and plant growth promoting traits of culturable endophytic actinobacteria associated with *Jatropha curcas* L. growing in Panxi dry-hot valley soil. *Appl Soil Ecol* 2015;93:47–55.
- [68] Madhaiyan M, Alex THH, Te Ngoh S, Prithiviraj B, Ji L. Leaf-residing Methylobacterium species fix nitrogen and promote biomass and seed production in *Jatropha curcas*. *Biotechnol Biofuels* 2015;8:1.
- [69] Mohanty SR, Dubey G, Kollah B. Endophytes of *Jatropha curcas* promote growth of maize. *Rhizosphere* 2016;3:20–8.
- [70] Jha CK, Patel D, Rajendran N, Saraf M. Combinatorial assessment on dominance and informative diversity of PGPR from rhizosphere of *Jatropha curcas* L. *J Basic Microbiol* 2010;50:211–7.
- [71] Jha CK, Patel B, Saraf M. Stimulation of the growth of *Jatropha curcas* by the plant growth promoting bacterium Enterobacter cancerogenus MSA2. *World J Microbiol Biotechnol* 2012;28:891–9.
- [72] Pant R, Pandey P, Kotoky R. Rhizosphere mediated biodegradation of 1, 4-di-chlorobenzene by plant growth promoting rhizobacteria of *Jatropha curcas*. *Ecol Eng* 2016;94:50–6.
- [73] Patel D, Saraf M. Influence of soil ameliorants and microflora on induction of antioxidant enzymes and growth promotion of *Jatropha curcas* L. under saline condition. *Eur J Soil Biol* 2013;55:47–54.
- [74] Sumarsih S, Haryanto D. *Pseudomonas fluorescens* and *Pseudomonas putida* for promoting growth of *Jatropha curcas* seedling root. *J Trop Life Sci* 2012;2:53–7.
- [75] Desai S, Narayanaiah C, Kumari CK, Reddy MS, Gnanamanickam SS, Rao GR, Venkateswarlu B. Seed inoculation with *Bacillus* spp. improves seedling vigour in oil-seed plant *Jatropha curcas* L. *Biol Fertil Soils* 2007;44:229–34.
- [76] Osorio LRM, Salvador AFT, Jongschaap REE, Perez CAA, Sandoval JEB, Trindade LM, Visser RGF, van Loo EN. High level of molecular and phenotypic biodiversity in *Jatropha curcas* from Central America compared to Africa, Asia and South America. *BMC Plant Biol* 2014;14:77.
- [77] Basha SD, Francis G, Makkar HPS, Becker K, Sujatha M. A comparative study of biochemical traits and molecular markers for assessment of genetic relationships between *Jatropha curcas* L. germplasm from different countries. *Plant Sci* 2009;176:812–23.
- [78] de Azevedo Peixoto L, Laviola BG, Alves AA, Rosado TB, Bhering LL. Breeding *Jatropha curcas* by genomic selection: a pilot assessment of the accuracy of predictive models. *PLoS One* 2017;12:0173368.
- [79] Mazumdar P, Basu A, Paul A, Mahanta C, Sahoo L. Age and orientation of the cotyledonary leaf explants determine the efficiency of *de novo* plant regeneration and *Agrobacterium tumefaciens*-mediated transformation in *Jatropha curcas* L. *S Afr J Bot* 2010;76:337–44.
- [80] Misra P, Toppo DD, Mishra MK, Saema S, Singh G. *Agrobacterium tumefaciens*-mediated transformation protocol of *Jatropha curcas* L. using leaf and hypocotyl segments. *J Plant Biochem Biotechnol* 2012;21:128–33.
- [81] Maravi DK, Kumar S, Sharma PK, Kobayashi Y, Goud VV, Sakurai N, Koyama H, Sahoo L. Ectopic expression of AtDGAT1, encoding diacylglycerol O-acyl-transferase exclusively committed to TAG biosynthesis, enhances oil accumulation in seeds and leaves of *Jatropha*. *Biotechnol Biofuels* 2016;9:226.
- [82] Joshi M, Jha A, Mishra A, Jha B. Developing transgenic *Jatropha* using the *SbNHX1* gene from an extreme halophyte for cultivation in saline wasteland. *PLoS One* 2013;8:1136.
- [83] Ye J, Qu J, Mao HZ, Ma ZG, Rahman NE, Bai C, Chen W, Jiang SY, Ramachandran S, Chua NH. Engineering geminivirus resistance in *Jatropha curcas*. *Biotechnol Biofuels* 2014;7:1.
- [84] Jones MH, Csurhes S. Pest plant risk assessment physic nut (*Jatropha curcas*). *Dep Prim Ind Fish, State Qld* 2008;1:28.
- [85] Yi C, Zhang S, Liu X, Bui HT, Hong Y. Does epigenetic polymorphism contribute to phenotypic variances in *Jatropha curcas* L.? *BMC Plant Biol* 2010;10:259.
- [86] Wu J, Liu Y, Tang L, Zhang F, Chen F. A study on structural features in early flower development of *Jatropha curcas* L. and the classification of its inflorescences. *Afr J Agric Res* 2011;6:275–84.
- [87] Kaur K, Dhillon GPS, Gill RIS. Floral biology and breeding system of *Jatropha curcas* in north-western India. *J Trop Sci* 2011;23:4–9.
- [88] Alam NCN, Abdullah TL, Abdullah NA. Flowering and fruit set under Malaysian climate of *Jatropha curcas* L. *Am J Agric Biol Sci* 2011;6:142–7.
- [89] Ahon LE, Quenum F. Floral biology and hybridization potential of nine accessions of physic nut (*Jatropha curcas* L.) originating from three continents. *Tropicultura* 2012;30:193–8.
- [90] Makwana V, Shukla P, Robin P. GA application induces alteration in sex ratio and cell death in *Jatropha curcas*. *Plant Growth Regul* 2010;61:121–5.
- [91] Pan BZ, Xu ZF. Benzyladenine treatment significantly increases the seed yield of the biofuel plant *Jatropha curcas*. *J Plant Growth Regul* 2011;30:166–74.
- [92] Ghosh A, Chikara J, Chaudhary DR, Prakash AR, Boricha G, Zala A. Paclobutrazol arrests vegetative growth and unveils unexpressed yield potential of *Jatropha curcas*. *J Plant Growth Regul* 2010;29:307–15.
- [93] Xu G, Luo R, Yao Y. Paclobutrazol improved the reproductive growth and the quality of seed oil of *Jatropha curcas*. *J Plant Growth Regul* 2013;32:875–83.
- [94] Juan S, MaoSheng C, JiaLong L, LongJian N, ZengFu X. Effects of soil-applied paclobutrazol on the vegetative and reproductive growth of biofuel plant *Jatropha curcas*. *Plant Divers Resour* 2013;35:173–9.
- [95] Dasumiaty, Miftahudin, Triadiati, Hartana A, Pronowo D. Increasing hermaphrodite flowers using plant growth regulators in andromonoecious *Jatropha curcas*. *HAYATI J Biosci* 2014;21:111–20.
- [96] Pan BZ, Luo Y, Song L, Chen MS, Li JL, Xu ZF. Thiazuron increases fruit number

- in the biofuel plant *Jatropha curcas* by promoting pistil development. *Ind Crop Prod* 2016;81:202–10.
- [97] Li C, Luo L, Fu Q, Niu L, Xu ZF. Isolation and functional characterization of JcFT, a FLOWERING LOCUS T (FT) homologous gene from the biofuel plant *Jatropha curcas*. *BMC Plant Biol* 2014;14:125.
- [98] Tang M, Tao YB, Fu Q, Song Y, Niu L, Xu ZF. An ortholog of LEAFY in *Jatropha curcas* regulates flowering time and floral organ development. *Sci Rep* 2016;6.
- [99] Xu G, Huang J, Yang Y, Yao YA. Transcriptome analysis of flower sex differentiation in *Jatropha curcas* L. Using RNA sequencing. *PLoS One* 2016;11:0145613.
- [100] Vaknin Y. The significance of pollination services for biodiesel feedstocks, with special reference to *Jatropha curcas* L.: a review. *Bioenergy Res* 2012;5:32–40.
- [101] Sukarin W, Yamada Y, Sakaguchi S. Characteristics of physic nut, *Jatropha curcas* L. as a new biomass crop in the tropics. *Jpn Agric Res Q* 1987;20:302–3.
- [102] Raju AS, Ezradanam V. Pollination ecology and fruiting behaviour in a monoeious species, *Jatropha curcas* L. (Euphorbiaceae). *Curr Sci* 2002;83:1395–8.
- [103] Bhattacharya A, Datta K, Kumar SD. Floral biology, floral resources constraints and pollination limitation in *Jatropha curcas* L. *Pak J Biol Sci* 2005;8:456–60.
- [104] Dhillon RS, Hooda MS, Handa AK, Ahlawat KS, Kumar Y, Subha SH, Singh N. Clonal propagation and reproductive biology of *Jatropha curcas* L. *Indian J Agroforest* 2006;8:18–27.
- [105] Santos MJ, Machado IC, Lopes AV. Biologia reprodutiva de duas espécies de *Jatropha* L. (Euphorbiaceae) em Caatinga, Nordeste do Brasil. *Rev Bras De Botânica* 2005;28:361–73.
- [106] Chang-Wei L, Kun L, You C, Yong-Yu S. Floral display and breeding system of *Jatropha curcas* L. *Stud China* 2007;9:114–9.
- [107] Quin Y, Ping PD, Biao DZ, Liang W, Xiang SQ. Study on pollination biology of *Jatropha curcas* (Euphorbiaceae). *J South China Agric Univ* 2007;28:62–6.
- [108] Rianti P, Suryobroto B, Atmowidi T. Diversity and effectiveness of insect pollinators of *Jatropha curcas* L. (Euphorbiaceae). *J Biosci* 2010;17:38–42.
- [109] Rincón-Rabanales M, Vargas-López LI, Adriano-Anaya L, Vázquez-Ovando A, Salvador-Figueroa M, Ovando-Medina I. Reproductive biology of the biofuel plant *Jatropha curcas* in its center of origin. *PeerJ* 2016;4:1819.
- [110] Samra S, Samocha Y, Eisikowitch D, Vaknin Y. Can ants equal honeybees as effective pollinators of the energy crop *Jatropha curcas* L. under Mediterranean conditions? *Glob Change Biol Bioenergy* 2014;6:756–67.
- [111] Kumar V, Makkar HPS, Becker K. Detoxified *Jatropha curcas* kernel meal as a dietary protein source: growth performance, nutrient utilization and digestive enzymes in common carp (*Cyprinus carpio* L.) fingerlings. *Aquac Nutr* 2011;17:313–26.
- [112] Xiao J, Zhang H. Comparative evaluation of *Jatropha curcas* L. seed meals obtained by different methods of defatting on toxic, antinutritional and nutritive factors. *J Food Sci Technol* 2014;51:1126–32.
- [113] Mahanta N, Gupta A, Khare SK. Production of protease and lipase by solvent tolerant *Pseudomonas aeruginosa* PseA in solid-state fermentation using *Jatropha curcas* seed cake as substrate. *Bioresour Technol* 2008;99:1729–35.
- [114] Ilmi M, Hidayat C, Hastuti P, Heeres HJ, Van der Maarel MJEC. Utilisation of *Jatropha* press cake as substrate in biomass and lipase production from *Aspergillus niger* 6516 and *Rhizomucor miehei* CBS 360.62. *Biocatal Agric Biotechnol* 2017;9:103–7.
- [115] Kannoju B, Ganapathiwar S, Nunavath H, Sunkar B, Bhukya B. Plausible exploitation of *Jatropha* de-oiled seed cake for lipase and phytase production and simultaneous detoxification by *Candida parapsilosis* isolated from poultry garbage. *Bioresour Technol* 2017;225:215–24.
- [116] Saetae D, Suntornsuk W. Toxic compound, anti-Nutritional factors and functional properties of protein isolated from Detoxified *Jatropha curcas* seed cake. *Int J Mol Sci* 2011;12:66–77.
- [117] Lin J, Zhou X, Wang J, Jiang P, Tang K. Purification and characterization of curcin, a toxic lectin from the seed of *Jatropha curcas*. *Prep Biochem Biotech* 2010;40:107–18.
- [118] Patade VY, Khatri D, Kumar K, Grover A, Kumari M, Gupta SM, Kumar D, Nasim M. RNAi mediated curcin precursor gene silencing in *Jatropha* (*Jatropha curcas* L.). *Mol Bio Rep* 2014;41:4305–12.
- [119] Lin J, Chen Y, Xu Y, Yan F, Tang L, Chen F. Cloning and expression of curcin, a ribosome-inactivating protein from the seeds of *Jatropha curcas*. *Acta Bot Sin* 2003;45(7).
- [120] Goel G, Makkar HP, Francis G, Becker K. Phorbol esters: structure, biological activity, and toxicity in animals. *Int J Toxicol* 2007;26:279–88.
- [121] Lin J, Chen Y, Xu Y, Yan F, Tang L, Chen F. Cloning and expression of curcin, a ribosome-inactivating protein from the seeds of *Jatropha curcas*. *Acta Bot Sin* 2003;45.
- [122] Adebawale AA, Kareem ST, Sobukola OP, Adebisi MA, Obadina AO, Kajihaua OE, Adegunwa MO, Sanni LO, Keith T. Mineral and Antinutrient content of high quality cassava-Tigernut composite flour extruded Snack. *J Food Process Pres* 2016.
- [123] Obasi NL, Ejikeme MP, Egbuonu CAC. Antimicrobial and phytochemical activity of methanolic extract and its fractions of *Jatropha curcas* Linn. (Euphorbiaceae) stem bark. *Afr J Pure Appl Chem* 2011;5:92–6.
- [124] Aregheore EM, Becker K, Makkar HPS. Detoxification of a toxic variety of *Jatropha curcas* using heat and chemical treatments, and preliminary nutritional evaluation with rats. *South Pac J Nat Appl Sci* 2003;21:51–6.
- [125] Makkar HP, Francis G, Becker K. Protein concentrate from *Jatropha curcas* screw-pressed seed cake and toxic and antinutritional factors in protein concentrate. *J Sci Food Agric* 2008;88:1542–8.
- [126] Harter T, Buhrke F, Kumar V, Focken U, Makkar HPS, Becker K. Substitution of fish meal by *Jatropha curcas* kernel meal: effects on growth performance and body composition of white leg shrimp (*Litopenaeus vannamei*). *Aquacult Nutr* 2011;17:542–8.
- [127] Wang WG, Li R, Liu B, Li L, Wang SH, Chen F. Effects of low nitrogen and drought stresses on proline synthesis of *Jatropha curcas* seedling. *Acta Physiol Plant* 2011;33:1591–5.
- [128] Brooker JD. Methods for detoxifying oil seed crops. United States Patent Application Publication. Appl. No. 13/133, 269. Publication.no.:US2011/0281017A1, 11 pp; 2011.
- [129] Pighinelli ALMT, Ferrari RA, Machado MCNA, Park KJ. Study of *Jatropha curcas* L. cake detoxification. Proceedings of the Energy, biomass and biological residues. In: International Conference of Agricultural Engineering – CIGR-AgEng 2012: Agriculture and Engineering for a Healthier Life 8–12 July 2012, Valencia, Spain, 2012, pp; 0340.
- [130] Rahma EH, Mansour EH, Hamoda ST. Biological evaluation of *Jatropha curcas* seed as a new source of protein. *Merit Res J Food Sci Technol* 2013;1:23–30.
- [131] Nokkaew R, Punsuvon V. Multistage solvent extraction for high yield oil and phorbol esters removal from Thai toxic *Jatropha curcas* meal. *Walailak J Sci Technol* 2015;12:299–301.
- [132] El Diwani GI, El Rafei SA, Hawash SI. Ozone for phorbol esters removal from Egyptian *Jatropha* oil seed cake. *Adv Appl Sci Res* 2011;2:221–32.
- [133] Makkar HP. State-of-the-art on detoxification of *Jatropha curcas* products aimed for use as animal and fish feed: a review. *Anim Feed Sci Technol* 2016;222:87–99.
- [134] Kasuya MCM, Montavani HC, da Luz JMR, da Silva JS, da Silva Pereira LP, Rodrigues MT. Bio-detoxification of *Jatropha* seed cake and its use in animal feed. INTECH Open Access Publisher; 2012.
- [135] Joshi C, Mathur P, Khare SK. Degradation of phorbol esters by *Pseudomonas aeruginosa* PseA during solid-state fermentation of deoiled *Jatropha curcas* seed cake. *Bioresour Technol* 2011;102:4815–9.
- [136] Oliveira ASD, Schwambach TI, +Sinhorin AP, Oliveira MRC, Alessi KC, Oliveira Filho FAD, Pina DDS. Capacity of ensilage of *Jatropha curcas* L. cake to degrade forbol esters. *Rev Bras Zootecn* 2012;41:1545–9.
- [137] Phengnum T, Suntornsuk W. Detoxification and anti-nutrients reduction of *Jatropha curcas* seed cake by *Bacillus* fermentation. *J Biosci Bioeng* 2013;115:168–72.
- [138] Gu K, Tian D, Mao H, Wu L, Yin Z. Development of marker-free transgenic *Jatropha curcas* producing curcin-deficient seeds through endospore-specific RNAi-mediated gene silencing. *BMC Plant Biol* 2015;15:242.
- [139] He W, King AJ, Khan MA, Cuevas JA, Ramaramanana D, Graham IA. Analysis of seed phorbol-ester and curcin content together with genetic diversity in multiple provenances of *Jatropha curcas* L. from Madagascar and Mexico. *Plant Physiol Biochem* 2011;49:1183–90.
- [140] Qin W, Huang M-X, Ying X, Zhang X-S, Fang C. Expression of a ribosome inactivating protein (curcin 2) in *Jatropha curcas* is induced by stress. *J Biosci* 2005;30:351–7.
- [141] Huang MX, Hou P, Wei Q, Xu Y, Chen F. A ribosome-inactivating protein (curcin 2) induced from *Jatropha curcas* can reduce viral and fungal infection in transgenic tobacco. *Plant Growth Regul* 2008;54:115–23.
- [142] Li C, Ng A, Xie L, Mao H, Qiu C, Srinivasan R, Yin Z, Hong Y. Engineering low phorbol ester *Jatropha curcas* seed by intercepting casbene biosynthesis. *Plant Cell Rep* 2016;35:103–14.
- [143] Natarajan P, Kanagasabapathy D, Gunadayalan G, Panchalingam J, Sugantham PA, Singh KK, Madasamy P. Gene discovery from *Jatropha curcas* by sequencing of ESTs from normalized and full-length enriched cDNA library from developing seeds. *BMC Genom* 2010;11:606.
- [144] Ohlrogge JB, Jaworski JG. Regulation of fatty acid synthesis. *Annu Rev Plant Biol* 1997;48:109–36.
- [145] Kennedy EP. Biosynthesis of complex lipids. *Fed Proc* 1961;20:934–40.
- [146] Slack CR, Campbell LC, Browse JA, Roughan PG. Some evidence for the reversibility of the cholinephosphotransferase-catalysed reaction in developing linseed cotyledons in vivo. *Biochim Biophys Acta* 1983;754:10–20.
- [147] Gu K, Chiam H, Tian D, Yin Z. Molecular cloning and expression of heteromeric ACCase subunit genes from *Jatropha curcas*. *Plant Sci* 2011;180:642–9.
- [148] Tong L, Shu-Ming P, Wu-Yuan D, Dan-Wei M, Ying X, Meng X, Fang C. Characterization of a new stearyl-acyl carrier protein desaturase gene from *Jatropha curcas*. *Biotechnol Lett* 2006;28:657–62.
- [149] Li J, Li MR, Wu PZ, Tian CE, Jiang HW, Wu GJ. Molecular cloning and expression analysis of a gene encoding a putative β -ketoacyl-acyl carrier protein (ACP) synthase III (KAS III) from *Jatropha curcas*. *Tree Physiol* 2008;28:921–7.
- [150] Wei Q, Li J, Zhang L, Wu P, Chen Y, Li M, Jiang H, Wu G. Cloning and characterization of a β -ketoacyl-acyl carrier protein synthase II from *Jatropha curcas*. *J Plant Physiol* 2012;169:816–24.
- [151] Wu PZ, Li J, Wei Q, Zeng L, Chen YP, Li MR, Jiang HW, Wu GJ. Cloning and functional characterization of an acyl-acyl carrier protein thioesterase (*JcFATB1*) from *Jatropha curcas*. *Tree Physiol* 2009;29:1299–305.
- [152] Wu P, Zhang S, Zhang L, Chen Y, Li M, Jiang H, Wu G. Functional characterization of two microsomal fatty acid desaturases from *Jatropha curcas* L. *J Plant Physiol* 2013;170:1360–6.
- [153] Nan Z, Ronghua X, Xiaozhu L, Aizhong L. Cloning and sequence analysis of sn-glycerol-3-phosphate acyltransferase gene (*JcGPAT*) from *Jatropha curcas*. *Plant Physiol Commun* 2011;47:181–8.
- [154] Misra A, Khan K, Niranjana A, Nath P, Sane VA. Over-expression of *JcDGAT1* from *Jatropha curcas* increases seed oil levels and alters oil quality in transgenic *Arabidopsis thaliana*. *Phytochem* 2013;96:37–45.
- [155] Xu R, Yang T, Wang R, Liu A. Characterisation of DGAT1 and DGAT2 from *Jatropha curcas* and their functions in storage lipid biosynthesis. *Funct Plant Biol* 2014;41:321–9.
- [156] Xu RH, Qiu LJ, Yang TQ, Wang RL, Tian B, Li AZ. Cloning and functioning of

- phospholipids: diacylglycerol acyltransferase (*JcPDAT1*) cDNA from *Jatropha curcas*. *Chin J oil. Crop Sci* 2013;35:123–30.
- [157] Palash SM, Kalam MA, Masjuki HH, Arbab MI, Masum BM, Sanjid A. Impacts of NO_x reducing antioxidant additive on performance and emissions of a multi-cylinder diesel engine fueled with *Jatropha* biodiesel blends. *Energ Convers Manag* 2014;77:577–85.
- [158] Hoekman SK, Broch A, Robbins C, Cenicerio E, Natarajan M. Review of biodiesel composition, properties, and specifications. *Renew Sustain Energ Rev* 2012;16:143–69.
- [159] Yaakob Z, Narayanan BN, Padikkaparambil S. A review on the oxidation stability of biodiesel. *Renew Sustain Energ Rev* 2014;35:136–53.
- [160] Yu N, Xiao WF, Zhu J, Chen XY, Peng CC. The *Jatropha curcas* KASIII gene alters fatty acid composition of seeds in *Arabidopsis thaliana*. *Biol Plant* 2015;59:773–82.
- [161] Qu J, Mao HZ, Chen W, Gao SQ, Bai YN, Sun YW, Geng YF, Ye J. Development of marker-free transgenic *Jatropha* plants with increased levels of seed oleic acid. *Biotechnol Biofuels* 2012;5:1.
- [162] Kim MJ, Yang SW, Mao H-Z, Veenia SP, Yin J-L, Chua N-H. Gene silencing of Sugar-dependent 1 (*JcSDP1*), encoding a patatin-domain triacylglycerol lipase, enhances seed oil accumulation in *Jatropha curcas*. *Biotechnol Biofuels* 2014;7:36.
- [163] Costa GG, Cardoso KC, Del Bem LE, Lima AC, Cunha MA, de Campos-Leite L, Vicentini R, Papes F, Moreira RC, Yunes JA, Campos FA. Transcriptome analysis of the oil-rich seed of the bioenergy crop *Jatropha curcas* L. *BMC Genom* 2010;11:462.
- [164] King AJ, Li Y, Graham IA. Profiling the developing *Jatropha curcas* L. seed transcriptome by pyrosequencing. *Bioenergy Res* 2011;4:211–21.
- [165] Silva-Junior OB, Rosado TB, Laviola BG, Pappas MR, Pappas GJ, Grattapaglia D. September. Genome-wide SNP discovery from a pooled sample of accessions of the biofuel plant *Jatropha curcas* based on whole-transcriptome Illumina resequencing. *BMC Proc* 2011;5:1.
- [166] Grover A, Kumari M, Singh S, Rathode SS, Gupta SM, Pandey P, Gilotra S, Kumar D, Arif M, Ahmed Z. Analysis of *Jatropha curcas* transcriptome for oil enhancement and genic markers. *Physiol Mol Biol Plants* 2014;20:139–42.
- [167] Gao S, Ouyang C, Wang S, Xu Y, Tang L, Chen F. Effects of salt stress on growth, antioxidant enzyme and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedlings. *Plant Soil Environ* 2008;54:374–81.
- [168] Niu G, Rodriguez D, Mendoza M, Jifon J, Ganjegunte G. Responses of *Jatropha curcas* to salt and drought stresses. *Int J Agron* 2012.
- [169] Dou XY, Wu GJ, Huang HY, Hou YJ, Gu Q, Peng CL. Responses of *Jatropha curcas* L. seedlings to drought stress. *Ying yong sheng tai xue bao* = *J Appl Ecol* 2008;19:1425–30.
- [170] Sapeta H, Costa JM, Lourenco T, Maroco J, Van der Linde P, Oliveira MM. Drought stress response in *Jatropha curcas*: growth and physiology. *J Exp Bot* 2013;85:76–84.
- [171] Gimeno V, Syvertsen JP, Simón I, Nieves M, Díaz-López L, Martínez V, García-Sánchez F. Physiological and morphological responses to flooding with fresh or saline water in *Jatropha curcas*. *Environ Exp Bot* 2012;78:47–55.
- [172] Tang M, Liu X, Deng H, Shen S. Over-expression of *JcDREB*, a putative AP2/EREBP domain-containing transcription factor gene in woody biodiesel plant *Jatropha curcas*, enhances salt and freezing tolerance in transgenic *Arabidopsis thaliana*. *Plant Sci* 2011;181:623–31.
- [173] Peng X, Liu H, Wang D, Shen S. Genome-wide identification of the *Jatropha curcas* MYB family and functional analysis of the abiotic stress responsive gene *JcMYB2*. *BMC Genom* 2016;17:1.
- [174] Tsuchimoto S, Cartagena J, Khemkladngoen N, Singkaravanit S, Kohinata T, Wada N, Sakai H, Morishita Y, Suzuki H, Shibata D, Fukui K. Development of transgenic plants in *Jatropha* with drought tolerance. *Plant Biotechnol* 2012;29:137–43.
- [175] Zhang L, Zhang C, Wu P, Chen Y, Li M, Jiang H, Wu G. Global analysis of gene expression profiles in physic nut (*Jatropha curcas* L.) seedlings exposed to salt stress. *PLoS One* 2014;9:97878.
- [176] Zhang C, Zhang L, Zhang S, Zhu S, Wu P, Chen Y, Li M, Jiang H, Wu G. Global analysis of gene expression profiles in physic nut (*Jatropha curcas* L.) seedlings exposed to drought stress. *BMC Plant Biol* 2015;15:1.
- [177] Wang H, Zou Z, Wang S, Gong M. Global analysis of transcriptome responses and gene expression profiles to cold stress of *Jatropha curcas* L. *PLoS One* 2013;8:82817.
- [178] Juntawong P, Sirikhachornkit A, Pimjan R, Sonthirod C, Sangsarakru D, Yoocha T, Tangphatsornruang S, Srinives P. Elucidation of the molecular responses to water-logging in *Jatropha* roots by transcriptome profiling. *Front Plant Sci* 2014;5:658.
- [179] Sharma S, Verma M, Sharma A. Utilization of non-edible oil seed cakes as substrate for growth of *Paecilomyces lilacinus* and as biopesticide against termites. *Waste Biomass- Valor* 2013;4:325–30.
- [180] Yun X, Huang Q, Rao W, Xiao C, Zhang T, Mao Z, Wan Z. A comparative assessment of cytotoxicity of commonly used agricultural insecticides to human and insect cells. *Ecotoxicol Environ Saf* 2017;137:179–85.
- [181] Vidau C, Diogon M, Aufaure J, Fontbonne R, Viguès B, Brunet JL, Texier C, Biron DG, Blot N, El Alaoui H, Belzunces LP. Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*. *PLoS One* 2011;6:21501.
- [182] Wang Z, Yan H, Yang Y, Wu Y. Biotype and insecticide resistance status of the whitefly *Bemisia tabaci* from China. *Pest Manag Sci* 2010;66:1360–6.
- [183] Djimilly YW, Nacro S. Natural enemies of *Calidea panaethiopica* (Heteroptera: Scutelleridae): an insect pest of *Jatropha curcas* L. in the South-Sudan zone of Burkina Faso. *Adv Entomol* 2016;4:201.
- [184] Schünemann R, Knaak N, Fiuza LM. Mode of action and specificity of *Bacillus thuringiensis* toxins in the control of caterpillars and stink bugs in soybean culture. *ISRN Microbiol* 2014;12.
- [185] Ibrahim MA, Griko N, Junker M, Bulla LA. *Bacillus thuringiensis*: a genomics and proteomics perspective. *Bioeng Bugs* 2010;1:31–50.
- [186] Gindin G, Navon A, Saphir N, Protasov A, Mendel Z. Environmental persistence of *Bacillus thuringiensis* products tested under natural conditions against *Thaumetopoea wilkinsoni*. *Phytoparasitica* 2007;35:255–63.
- [187] Mazumdar P. Germplasm Evaluation, Environmental Impact Assessment and Genetic Improvement Studies in *Jatropha curcas* (Doctoral dissertation); 2012.
- [188] Gu K, Mao H, Yin Z. Production of marker-free transgenic *Jatropha curcas* expressing hybrid *Bacillus thuringiensis* δ -endotoxin Cry1Ab/1Ac for resistance to larvae of tortrix moth (*Archips micaceanus*). *Biotechnol Biofuels* 2014;7:68.
- [189] QiaoSheng G, YuChuan L, QingSong S, Fei G, XueHai Z. Bioactivity of extracts from *Jatropha curcas* L. branches and leaves against plant pathogenic fungi. *Bull Bot Res Harbin* 2010;30:503–7.
- [190] Hu H-R, Sun Y-C, Chen F, Sun Q. Pathogen identification of *Jatropha curcas* L. wilt disease and screening of its fungicides. *J Sichuan Agric Univ* 2009;6:047.
- [191] Kumar S, Sharma S, Pathak DV, Beniwal J. Integrated management of *Jatropha* root rot caused by *Rhizoctonia bataticola*. *J Trop Sci* 2011:35–41.
- [192] Shimizu T, Nakazono-Nagaoka E, Uehara-Ichiki T, Sasaya T, Omura T. Targeting specific genes for RNA interference is crucial to the development of strong resistance to Rice stripe virus. *Plant Biotechnol J* 2011;9:503–12.
- [193] Tenllado F, Martínez-García B, Vargas M, Díaz-Ruiz JR. Crude extracts of bacterially expressed dsRNA can be used to protect plants against virus infections. *BMC Biotechnol* 2003;3:3.
- [194] Lau SE, Mazumdar P, Hee TW, Song ALA, Othman RY, Hari Krishna JA. Crude extracts of bacterially-expressed dsRNA protect orchid plants against Cymbidium mosaic virus during transplantation from in vitro culture. *J Hort Sci Biotechnol* 2014;89:569–76.
- [195] Mitter N, Worrall EA, Robinson KE, Li P, Jain RG, Taochy C, Fletcher SJ, Carroll BJ, Lu GQ, Xu ZP. Clay nanosheets for topical delivery of RNAi for sustained protection against plant viruses. *Nat Plants* 2017;3. [16207-16207].
- [196] Georges F, Ray H. Genome Editing of Crops: a Renewed Opportunity for Food Security. *GM Crops Food* 2017;8:1–12.
- [197] Woo JW, Kim J, Kwon SI, Corvalán C, Cho SW, Kim H, Kim SG, Kim ST, Choe S, Kim JS. DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat Biotechnol* 2015;33:1162–4.
- [198] de Barros CR, Ferreira LM, Nunes FM, Bezerra RM, Dias AA, Guedes CV, Cone JW, Marques GS, Rodrigues MA. The potential of white-rot fungi to degrade phorbol esters of *Jatropha curcas* L. seed cake. *Eng Life Sci* 2011;11:107–10.
- [199] Belew MA, Belew KY, Lawal IA. Cocktail of fungi blend on *Jatropha curcas* kernel cake: Effect on feed intake and blood parameters of goat. *Libyan Agric Res J* 2011;2:138–43.
- [200] Sharath BS, Mohankumar BV, Somashekar D. Bio-detoxification of phorbol esters and other anti-nutrients of *Jatropha curcas* seed cake by fungal cultures using solid-state fermentation. *Appl Biochem Biotechnol* 2014;172:2747–57.
- [201] da Luz JEMR, Nunes MD, Paes SA, Torres DP, Kasuya MCM. Bio-detoxification of *Jatropha curcas* seed cake by *Pleurotus ostreatus*. *Afr J Microbiol Res* 2014;8:1148–56.
- [202] Foidl N, Foidl G, Sanchez M, Mittelbach M, Hackel S. *Jatropha curcas* L. as a source for the production of biofuel in Nicaragua. *Bioresour Technol* 1996;58:77–82.
- [203] Rashid U, Anwar F, Jamil A, Bhatti HN. *Jatropha curcas* seed oil as a viable source for biodiesel. *Pak J Bot* 2010;42:575–82.
- [204] Lu H, Liu Y, Zhou H, Yang Y, Chen M, Liang B. Production of biodiesel from *Jatropha curcas* L. oil. *Comput Chem Eng* 2009;33:1091–6.
- [205] Akintayo ET. Characteristics and composition of *Parkia biglobbosa* and *Jatropha curcas* oils and cakes. *Bioresour Technol* 2004;92:307–10.
- [206] Akbar E, Yaakob Z, Kamarudin SK, Ismail M, Salimon J. Characteristic and composition of *Jatropha curcas* oil seed from Malaysia and its potential as biodiesel feedstock feedstock. *Eur J Sci Res* 2009;29:396–403.
- [207] Belew M, Adekola F, Adebayo G, Ameen O, Muhammed N, Olaniyan A, Adekola O, Musa A. Physico-chemical characteristics of oil and biodiesel from Nigerian and Indian *Jatropha curcas* seeds. *IJBCT* 2010:4.
- [208] de Oliveira JS, Leite PM, de Souza LB, Mello VM, Silva EC, Rubim JC, Meneghetti SM, Suarez PA. Characteristics and composition of *Jatropha gossypifolia* and *Jatropha curcas* L. oils and application for biodiesel production. *Biomass- Bioenerg* 2009;33:449–53.
- [209] Qian J, Shi H, Yun Z. Preparation of biodiesel from *Jatropha curcas* L. oil produced by two-phase solvent extraction. *Bioresour Technol* 2010;101:7025–31.
- [210] Deng X, Fang Z, Liu YH, Yu CL. Production of biodiesel from *Jatropha* oil catalyzed by nanosized solid basic catalyst. *Energy* 2011;36:777–84.
- [211] Ojolo SJ, Adelaja AO, Ogbonnaya M, Ogunsin BS. Study of an effective technique for the production of biodiesel from *Jatropha* oil. *J Emerg Trends Eng Appl Sci* 2011;2:79–86.
- [212] Bueso F, Moreno L, Cedeño M, Manzanarez K. Lipase-catalyzed biodiesel production and quality with *Jatropha curcas* oil: exploring its potential for Central America. *J Biol Eng* 2015;9:12.
- [213] Nalgundwar A, Paul B, Sharma SK. Comparison of performance and emissions characteristics of DI CI engine fuelled with dual biodiesel blends of palm and *Jatropha*. *Fuel* 2016;173:172–9.
- [214] Meshram PB, Joshi KC. A new report of *Spodoptera litura* (Fab.) Boursin (Lepidoptera: Noctuidae) as a pest of *Jatropha curcas* Linn. *Indian For* 1994;120:273–4.
- [215] Kumar A, Singh N. First report of *Maconellicoccus hirsutus* green infestation on *Jatropha curcas* saplings. *Phytoparasitica* 2014;42:71–3.
- [216] Arif M, Sharma S, Das SC. Incidence of leaf miner on *Jatropha curcas*-A bio diesel plant in secunderabad, India. *J Exp Zool* 2007;10:107.

- [217] Shanker C, Dhyani SK. Insect pests of *Jatropha curcas* L. and the potential for their management. *Curr Sci* 2006;91:162–3.
- [218] Sajap AS. First report of a field outbreak of a leaf miner, *Stomphastis thraustica* meyrick (Lepidoptera: Gracillariidae) on bioenergy crop *Jatropha curcas* L. (Euphorbiaceae) in Malaysia. *Serangga* 2010;15:65–71.
- [219] YinBo X, JianHua Z, Yan L, DeKui Z, Bo F. Morphological and biological observations on *Stomphastis thraustica* Meyrick (Lepidoptera: Gracillariidae), a leaf miner of *Jatropha curcas*. *Acta Entomol Sin* 2009;52:228–33.
- [220] Terren M, Mignon J, De Clerck C, Jijakli H, Savery S, Jacquet de Haveskercke P, Winandy S, Mergeai G. Principal disease and insect pests of *Jatropha curcas* L. in the lower Valley of the Senegal River. *Tropicultura* 2012;30:222–9.
- [221] Habou ZA, Adam T, Haubruge E, Mergeai G, Verheggen FJ. Insects associated with *Jatropha curcas* Linn. (Euphorbiaceae) in West Niger. *J Insect Sci* 2014;14:255.
- [222] Kumar R, Sinha A, Singh SR, Kamil D. Incidence of a leaf spot disease in *Jatropha curcas* from eastern Uttar Pradesh. *J Mycol Plant Pathol* 2009;39:536.
- [223] Sharma S, Kaushik JC, Kaushik N. *Fusarium moniliforme* causing root rot of *Jatropha*. *Indian Phytopathol* 2001;54:275.
- [224] Ferreira-Pinto MM, Silva MJ, Santos MR. Screening of *Jatropha curcas* genotypes to anthracnose caused by *Colletotrichum gloeosporioides*. *Commun Agric Appl Biol Sci* 2010;76:629–34.
- [225] Tewari JP, Dwivedi HD, Pathak M, Srivastava SK. Incidence of a mosaic disease in *Jatropha curcas* L. from eastern Uttar Pradesh. *Curr Sci* 2007;93:1048–9.
- [226] Durrett TP, Benning C, Ohlrogge J. Plant triacylglycerols as feedstocks for the production of biofuels. *Plant J* 2008;54:593–607.
- [227] Sood A, Chauhan RS. Regulation of FA and TAG biosynthesis pathway genes in endosperms and embryos of high and low oil content genotypes of *Jatropha curcas* L. *Plant Physiol Biochem* 2015;94:253–67.